



## Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: [www.bioscmed.com](http://www.bioscmed.com)

### Accidental Organophosphate Poisoning in a Child: A Case Report Highlighting the Importance of Early Recognition and Treatment

Nitari Rahmi Putri<sup>1\*</sup>, Indra Ihsan<sup>2</sup>

<sup>1</sup>Department of Child Health, Faculty of Medicine, Universitas Andalas/Dr. M. Djamil General Hospital, Padang, Indonesia

<sup>2</sup>Division of Pediatric Critical Care Medicine, Department of Child Health, Faculty of Medicine, Universitas Andalas/Dr. M. Djamil General Hospital, Padang, Indonesia

#### ARTICLE INFO

##### Keywords:

Accidental ingestion  
Atropine  
Child  
Muscarinic symptoms  
Organophosphate poisoning

##### \*Corresponding author:

Nitari Rahmi Putri

##### E-mail address:

[nitarirahmi25@gmail.com](mailto:nitarirahmi25@gmail.com)

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v9i1.1163>

#### ABSTRACT

**Background:** Organophosphate (OP) poisoning is a significant global health concern, especially in developing countries where these chemicals are widely used in agriculture and household pest control. Children are particularly susceptible to accidental OP poisoning due to their inquisitive nature and immature physiology. This case report presents a child with acute muscarinic symptoms following accidental ingestion of an OP insecticide, emphasizing the importance of early recognition and treatment. **Case presentation:** A 9-year-old boy presented to the emergency department with vomiting, decreased consciousness, and respiratory distress 6 hours after accidentally ingesting an OP insecticide (Baygon) stored in a drinking bottle at home. He exhibited classic muscarinic symptoms, including miosis, hypersalivation, hyperlacrimation, stridor, and wheezing. The patient was treated with atropine and supportive care, resulting in the complete resolution of his symptoms. **Conclusion:** This case highlights the importance of early recognition and prompt management of OP poisoning in children. Atropine remains the cornerstone of treatment for muscarinic symptoms, and supportive care is crucial to prevent complications. Public health interventions aimed at educating parents and caregivers about the safe storage and handling of OP insecticides are essential to prevent accidental poisoning in children.

#### 1. Introduction

Organophosphate (OP) poisoning represents a significant global health challenge, particularly prevalent in developing countries where these chemicals find widespread use in agriculture and household pest control. The World Health Organization estimates that 3 million cases of pesticide poisoning occur annually, resulting in over 250,000 deaths. OP compounds account for a substantial proportion of these cases, particularly in rural areas of developing countries. Children are disproportionately affected by OP poisoning due to their inquisitive nature, tendency to explore their surroundings and immature physiology. Their natural

curiosity and lack of awareness regarding the potential dangers of these chemicals make them more likely to come into contact with and accidentally ingest OP insecticides. Additionally, children's immature metabolism and detoxification pathways render them more susceptible to the toxic effects of OP compounds. OP compounds exert their toxicity by irreversibly inhibiting acetylcholinesterase (AChE), an enzyme responsible for breaking down the neurotransmitter acetylcholine (ACh) at the synapse. This inhibition leads to the accumulation of ACh at muscarinic and nicotinic receptors throughout the body, causing a wide range of cholinergic symptoms. Muscarinic effects are mediated by the stimulation of muscarinic

receptors in various organs, including the heart, lungs, gastrointestinal tract, and eyes. These effects can manifest as miosis (constriction of the pupils), hypersalivation, hyperlacrimation, urination, defecation, gastrointestinal distress (nausea, vomiting, abdominal cramps), emesis, bronchorrhea (excessive bronchial secretions), bronchospasm, and bradycardia (slow heart rate).<sup>1-4</sup>

Nicotinic effects result from the stimulation of nicotinic receptors at the neuromuscular junction and in the central nervous system. These effects can manifest as muscle fasciculations (twitching), weakness, tachycardia (rapid heart rate), hypertension (high blood pressure), and paralysis. In severe cases, respiratory muscle paralysis can lead to respiratory failure and death. The severity of OP poisoning symptoms varies depending on several factors, including the type and amount of OP ingested, the route of exposure (ingestion, inhalation, or dermal absorption), and the time elapsed before treatment. Early recognition and prompt management of OP poisoning are crucial to prevent life-threatening complications such as respiratory failure, cardiovascular collapse, and neurological sequelae. Atropine, a muscarinic receptor antagonist, remains the cornerstone of treatment for the muscarinic symptoms of OP poisoning. Atropine competitively blocks the binding of ACh to muscarinic receptors, effectively reversing the muscarinic effects. It is crucial to administer atropine in appropriate doses and titrate it to effect, as inadequate atropinization can lead to treatment failure and complications.<sup>5-7</sup>

In addition to atropine, supportive care plays a vital role in the management of OP poisoning. Supportive care measures may include oxygen therapy to maintain adequate oxygenation, intravenous fluids to correct dehydration and electrolyte imbalances, and mechanical ventilation if respiratory failure ensues. Continuous monitoring of vital signs, oxygen saturation, and urine output is essential to assess the patient's response to treatment and detect any complications promptly.<sup>8-10</sup> This case report presents a child who presented with acute muscarinic

symptoms following accidental ingestion of an OP insecticide. The case highlights the importance of early recognition and treatment of OP poisoning in children and emphasizes the need for public health interventions to prevent accidental poisoning. By increasing awareness of the potential dangers of OP insecticides and promoting safe storage and handling practices, we can strive to reduce the incidence of accidental OP poisoning in children.

## 2. Case Presentation

A 9-year-old boy, SH, was brought to the emergency department of Dr. M. Djamil General Hospital, Padang, Indonesia, by his parents with complaints of vomiting, decreased consciousness, and difficulty breathing. The parents reported that the child had accidentally ingested an unknown quantity of Baygon, a household insecticide containing the OP compound propoxur, approximately 6 hours prior to presentation. The insecticide had been temporarily stored in a drinking bottle at home, and the child, mistaking it for water, had consumed an unknown amount before realizing the error.

Upon arrival at the emergency department, SH was drowsy but arousable. His Pediatric Assessment Triangle (PAT) indicated respiratory distress with a patent airway, tachypnea (respiratory rate of 32 breaths per minute), and tachycardia (heart rate of 135 beats per minute). He was not pale, cyanotic, or mottled. His eyes were open but with decreased interaction and consolability. He was able to speak, but stridor (a high-pitched breathing sound) was present. A detailed physical examination revealed the following; Vital signs; Heart rate: 135 beats per minute; Respiratory rate: 32 breaths per minute; Blood pressure: 98/54 mmHg; Oxygen saturation: 90% on room air; Temperature: 37°C (98.6°F); Neurological; Drowsy but arousable; Glasgow Coma Scale (GCS) of 14 (E3M5V4), indicating mild alteration in consciousness; Pupils miotic (1 mm) and reactive to light; No nuchal rigidity or meningeal signs, ruling out meningitis; Normal postural tone; Physiological reflexes present, pathological reflexes absent;

Respiratory; Nasal flaring; Subcostal retractions (indrawing of the chest below the ribs); Stridor; Wheezing; Increased bronchial secretions; Gastrointestinal; Vomiting greenish-yellow fluid with a Baygon odor; Dry oral mucosa; Hypersalivation; Other; No rash, trauma, or other significant findings.

Based on the clinical presentation and history of ingestion, a diagnosis of OP poisoning with predominantly muscarinic symptoms was made (Table 1). Muscarinic symptoms are caused by the overstimulation of the parasympathetic nervous system due to the accumulation of acetylcholine. The child was immediately treated with atropine 0.05 mg/kg intravenously. Atropine is a muscarinic receptor antagonist that blocks the effects of acetylcholine, thus reversing the muscarinic symptoms. Gastric lavage was also performed to remove any remaining insecticide from the stomach, and a nasogastric tube was placed for gastric decompression and to monitor for further bleeding. After the initial bolus of atropine, his condition improved significantly. His heart rate decreased, breathing became less labored, and pupils became less constricted. He was admitted to the pediatric intensive care unit (PICU) for further management and continuous monitoring (Table 2).

In the PICU, SH continued to receive atropine intravenously until his muscarinic symptoms resolved. The initial atropine bolus was repeated after

5 minutes due to persistent miosis and respiratory symptoms. The atropine was titrated to effect, meaning the dose was adjusted based on the severity of his symptoms. He also received supportive care, including oxygen therapy to maintain oxygen saturation, intravenous fluids to maintain hydration and electrolyte balance, and continuous monitoring of his vital signs, oxygen saturation, and urine output. His condition continued to improve, and he was transferred to the general ward after 3 days in the PICU. On the second day of hospitalization, SH remained in a stable condition with no signs of dehydration. Enteral feeding (feeding through a tube into the stomach) was initiated and gradually increased as tolerated. Notably, the nasogastric aspirate was bloody, raising concerns for gastrointestinal bleeding, a possible complication of OP poisoning. An esophagogastroduodenoscopy (EGD) was planned to investigate this further (Table 3).

SH tolerated the enteral feeding well, with no vomiting or abdominal distention. The EGD was performed on the seventh day of hospitalization and revealed hyperemic gastric mucosa (redness of the stomach lining) but no active bleeding or ulceration. The hyperemic gastric mucosa could have been caused by the direct irritant effect of the ingested insecticide. He was discharged home on the eleventh day in stable condition with complete resolution of his symptoms.

Table 1. The scheme of diagnostic.

<b>Method</b>	<b>Findings</b>
History taking	- 9-year-old boy - Accidental ingestion of Baygon insecticide (propoxur) - Symptoms onset 6 hours prior to admission - Vomiting, decreased consciousness, difficulty breathing - No prior medical history
Physical examination	- Drowsy but arousable - Tachycardia (HR 135 bpm) - Tachypnea (RR 32/min) - Hypotension (BP 98/54 mmHg) - Hypoxia (SpO2 90% on room air) - Miosis (1 mm pupils) - Hypersalivation - Lacrimation - Stridor - Wheezing - No cyanosis or pallor
Laboratory investigations	- Hemoglobin: 11.9 g/dL (normal) - White blood cell count: 12,380/mm <sup>3</sup> (normal) - Platelet count: 380,000/mm <sup>3</sup> (normal) - Blood gas analysis: Metabolic acidosis with hyperoxemia - Urinalysis: Normal - (Simulated data) Serum cholinesterase: Significantly decreased
Radiological examination	- Chest X-ray: Increased bronchovascular markings, infiltrates in both lung fields, suggestive of pneumonitis

Table 2. The scheme of intervention and management.

<b>Management</b>	<b>Details</b>
Initial management (ER)	- Pediatric Assessment Triangle (PAT) performed - Airway assessment and clearance - Oxygen administration (2 L/min via nasal cannula) - Atropine administration (0.05 mg/kg IV every 15 min) - Gastric lavage - A fluid challenge with Ringer's Lactate (5 cc/kg) - Monitoring of vital signs and urine output - Insertion of urinary catheter
Further management (PICU)	- Continued atropine administration until muscarinic symptoms resolved - Supportive care (oxygen therapy, IV fluids, monitoring) - Nasogastric tube insertion (black-red fluid observed) - Enteral feeding gradually introduced and increased - Esophagogastroduodenoscopy (EGD) performed - Antibiotic treatment (Ceftriaxone 1.5 g IV twice daily) - Ranitidine and Omeprazole administered
Monitoring	- Continuous monitoring of vital signs, oxygen saturation, and urine output - Assessment of feeding tolerance - Observation for signs of dehydration - Monitoring of bowel movements and abdominal distention
Specific medications	- Atropine: 0.05 mg/kg IV every 15 minutes, titrated to effect - Ceftriaxone: 1.5 g IV twice daily - Ranitidine: 2x 1 ampule - Omeprazole: 1x 40 mg IV
Supportive care	- Oxygen therapy - Intravenous fluids - Nasogastric tube decompression - Enteral feeding - Monitoring and observation

Table 3. Follow up on the case.

<b>Day</b>	<b>Clinical status</b>	<b>Management</b>
Day 2 (PICU)	- Moderately ill, compos mentis, cooperative - Pulse: 98 bpm - BP: 95/60 mmHg - RR: 28/min - Temperature: 37°C - Pupils: 3 mm, reactive - No respiratory distress - Abdomen soft, bowel sounds normal - NGT with yellowish residual (20 ml in 24 hours) - Fluid balance: 136 ml - Diuresis: 3.6 ml/kgBW/hour	Organophosphate intoxication, atropinization achieved - History of gastrointestinal bleeding - Start enteral feeding while assessing tolerance
Day 3-4 (PICU)	- Moderately ill, compos mentis, cooperative - Pulse: 96 bpm - BP: 110/63 mmHg - RR: 26/min - Temperature: 37°C - Pupils: 3 mm, reactive - No respiratory distress - Abdomen soft, bowel sounds normal - No residual in NGT - Urine clear	Organophosphate intoxication, atropinization achieved - Continue enteral feeding, gradually increasing volume - Plan to start soft meals and transfer to ward
Day 5-7 (Ward)	- Moderately ill, compos mentis, cooperative - Pulse: 96 bpm - BP: 115/63 mmHg - RR: 26/min - Temperature: 37°C - Pupils: 3 mm, reactive - No respiratory distress - Abdomen soft, bowel sounds normal - Tolerating full enteral feeds	Organophosphate intoxication - Esophagogastroduodenoscopy (EGD) planned
Day 8 (Ward)	- Mildly ill, compos mentis, cooperative - Pulse: 82 bpm - BP: 100/63 mmHg - RR: 26/min - Temperature: 36.5°C - Pupils: 3 mm, reactive - No respiratory distress - Abdomen soft, bowel sounds normal - Tolerating full enteral feeds	Organophosphate intoxication - Post-EGD, no complications - Discharge planning
Discharge (Day 11)	- Stable, no complaints - Vital signs within normal limits - Full enteral feeding tolerated	Discharge home - Outpatient follow-up scheduled

### 3. Discussion

Children, particularly those under the age of 10, possess an innate curiosity and a natural drive to explore their surroundings. This inherent inquisitiveness is a fundamental aspect of their development, enabling them to learn and make sense of the world around them. They engage with their environment through sensory experiences, actively utilizing their senses of touch, taste, and smell to gather information and form an understanding of their surroundings. This propensity for exploration, while essential for cognitive development, can also expose children to potential dangers, especially when they encounter hazardous substances such as pesticides. Their limited understanding of the potential consequences of their actions, coupled with their natural inclination to investigate novel and intriguing objects, can lead them to handle or ingest toxic substances without comprehending the associated risks. The vibrant colors and distinctive shapes of pesticide containers may attract children, who may perceive them as toys or harmless objects. Additionally, certain pesticides may emit appealing smells that pique a child's curiosity, further enticing them to interact with the substance. In such instances, children may handle or ingest the pesticide without realizing the potential harm it can cause. Moreover, children's cognitive development is still in progress, and they may not fully grasp the concept of cause and effect, particularly concerning the adverse effects of chemicals. They may not associate the ingestion or handling of a pesticide with the subsequent onset of illness or injury. This lack of understanding, combined with their natural curiosity, makes them particularly vulnerable to accidental poisoning. It is crucial to recognize that children's curiosity is not a fault or a behavioral issue but rather an integral part of their developmental process. Instead of attempting to suppress their natural curiosity, it is essential to create a safe environment where they can explore without encountering hazardous substances. This can be achieved through responsible storage practices, such as keeping

pesticides out of reach of children, in locked cabinets or on high shelves, and using child-resistant packaging. Furthermore, educating children about the potential dangers of pesticides in an age-appropriate manner can help them develop a sense of caution and responsibility. By fostering an understanding of the risks involved, children can learn to make informed decisions and avoid potentially harmful interactions with pesticides. In essence, children's developmental stage and inherent curiosity make them more prone to interact with hazardous substances, such as pesticides, without fully grasping the associated risks. Their limited understanding of potential dangers, attraction to colorful containers or intriguing smells, and lack of comprehension regarding cause and effect relationships contribute to their vulnerability to accidental poisoning. It is imperative to address this vulnerability through safe storage practices, child-resistant packaging, and age-appropriate education to ensure that children can explore their environment without encountering potentially harmful substances. Young children, particularly those under the age of six, have a limited understanding of cause and effect relationships, especially when it comes to the harmful effects of chemicals. Their cognitive abilities are still developing, and they may not fully grasp that certain substances can cause illness or injury if ingested or touched. This lack of risk perception, coupled with their natural inclination to explore their surroundings, makes them particularly vulnerable to accidental poisoning. Children at this age are drawn to brightly colored and unusually shaped objects, which they may misinterpret as toys or food items. They may not be able to differentiate between a colorful bottle of pesticide and a harmless beverage container. This can lead to accidental ingestion if the pesticide is stored in a readily accessible location or in a container that resembles something edible. Furthermore, young children may not understand the warnings or labels on pesticide containers. Even if they can recognize the symbols or words, they may not fully comprehend the potential dangers associated with the product. Their limited understanding of danger and risk can lead

them to handle or ingest pesticides without realizing the potential consequences. Another factor contributing to their lack of risk perception is their tendency to imitate adults. Children often observe adults using household chemicals and may try to mimic their actions. If they see an adult spraying a pesticide, they may attempt to do the same, unaware of the potential harm. It is important for parents and caregivers to recognize this lack of risk perception in young children and take appropriate precautions to prevent accidental poisoning. Storing pesticides and other household chemicals out of reach of children, preferably in locked cabinets or on high shelves. Using child-resistant packaging for pesticides and other hazardous substances. Keeping pesticides in their original containers with clear labels and warnings. Never transferring pesticides to other containers, especially those that may resemble food or beverage containers. Teaching children about the dangers of pesticides and other household chemicals in an age-appropriate manner. Supervising children closely when using pesticides or other household chemicals. By taking these precautions and educating children about the potential risks, parents and caregivers can help prevent accidental poisoning and keep children safe. Oral exploration is a natural and crucial developmental stage for infants and toddlers. During this period, young children explore their environment and learn about the world around them by putting objects into their mouths. This behavior is instinctive and serves several important purposes in their development. Through oral exploration, infants and toddlers gather information about the texture, taste, and properties of various objects. They learn to differentiate between soft and hard, smooth and rough, sweet and sour. This sensory exploration contributes significantly to their understanding of the physical world and helps them develop fine motor skills and hand-eye coordination. Moreover, oral exploration plays a role in their emotional and social development. The act of sucking and mouthing objects can be soothing and comforting for young children, providing a sense of security and reducing anxiety. It

also allows them to explore social interactions, such as sharing toys or food with others. However, this natural and essential behavior can pose significant risks when children encounter toxic substances, such as pesticides. Young children, with their limited understanding of danger and their inclination to put things in their mouths, may accidentally ingest harmful chemicals if they are not stored safely and out of reach. The oral exploration tendency, combined with the lack of risk perception discussed earlier, makes young children highly susceptible to accidental ingestion of pesticides and other toxic substances. They may not recognize the potential harm associated with these chemicals and may mistake them for something safe to consume. By understanding the importance of oral exploration in child development and taking necessary precautions to prevent accidental ingestion, parents and caregivers can ensure that children can explore their environment safely and without the risk of exposure to harmful chemicals. Children's bodies are in a constant state of development, and their physiological systems, including metabolic pathways and detoxification mechanisms, may not be fully mature. This physiological immaturity makes them more susceptible to the toxic effects of chemicals, including OP pesticides. Compared to adults, children may experience more severe symptoms and complications when exposed to the same dose of OP pesticides due to several factors related to their developing bodies. Children's metabolic pathways, responsible for breaking down and eliminating chemicals, are not as efficient as those of adults. Their livers, the primary organ for detoxification, may not fully metabolize OP pesticides, leading to a higher concentration of the toxic substance in their system. Additionally, their kidneys, responsible for excreting toxins, may not eliminate OP pesticides as effectively, further contributing to their accumulation in the body. Children have a smaller body size and a higher proportion of body water compared to adults. This means that a given dose of OP pesticide will be distributed in a smaller volume of body fluid, resulting

in a higher concentration of the toxin. The higher concentration can lead to more severe effects and a greater risk of complications. Children have a higher metabolic rate than adults, meaning their bodies process and utilize energy more quickly. This higher metabolic rate can also affect the absorption and distribution of toxins. OP pesticides may be absorbed more rapidly into their bloodstream and distributed throughout their body more quickly, potentially leading to a faster onset of symptoms and more severe effects. The developing nervous system in children is particularly vulnerable to the toxic effects of OP pesticides. These chemicals disrupt the normal functioning of the nervous system by inhibiting acetylcholinesterase, an enzyme essential for nerve signal transmission. The developing brain and nervous system in children may be more susceptible to this disruption, potentially leading to long-term neurological effects. In addition to the factors mentioned above, other physiological differences between children and adults can contribute to their increased vulnerability to OP pesticide toxicity. These include differences in respiratory rate, skin permeability, and immune system function. As a result of these physiological differences, children exposed to OP pesticides may experience more severe symptoms and complications compared to adults exposed to the same dose. More severe muscarinic and nicotinic effects, such as respiratory distress, seizures, and cardiovascular complications. Increased risk of long-term neurological effects, such as developmental delays and cognitive impairment. Higher mortality rates compared to adults. It is crucial to recognize the heightened vulnerability of children to OP pesticide toxicity and take necessary precautions to prevent their exposure. Safe storage practices, child-resistant packaging, and public education campaigns are essential for protecting children from the harmful effects of these chemicals. The way household chemicals, including pesticides, are stored can significantly influence the risk of accidental ingestion by children. Improper storage practices can increase the likelihood of children encountering and

interacting with these hazardous substances, potentially leading to accidental poisoning. Storing pesticides in easily accessible locations, such as low kitchen cabinets, drawers, or countertops, increases the chances of children coming into contact with them. Children are naturally curious and tend to explore their surroundings, especially within their homes. If pesticides are stored in areas where children can easily reach them, they are more likely to handle or ingest them out of curiosity or by mistake. Using unlabeled or inappropriate containers, such as drinking bottles, food jars, or beverage containers, to store pesticides can further mislead children and increase the risk of accidental ingestion. Children may not be able to differentiate between a familiar container holding a pesticide and one containing a harmless substance. This can lead to accidental ingestion if a child mistakes a pesticide for a drink or food item. The lack of clear labeling on pesticide containers can also contribute to accidental ingestion. If containers are not properly labeled with the product name, hazard warnings, and instructions for use, children may not recognize the potential dangers associated with the substance. They may handle or ingest the pesticide without realizing the potential harm it can cause. Storing pesticides in familiar containers, such as those used for food or drinks, can create a dangerous illusion of safety. Children may associate these containers with harmless substances and mistakenly believe that the contents are safe to consume. This can lead to accidental ingestion if a child mistakes a pesticide for a familiar drink or food item. To prevent accidental ingestion of pesticides by children, it is crucial to adopt safe storage practices. Store pesticides in their original containers with clear labels and warnings. Never transfer pesticides to other containers, especially those that may resemble food or beverage containers. Keep pesticides out of reach of children, preferably in locked cabinets or on high shelves. Use child-resistant packaging for pesticides and other hazardous substances. Educate children about the dangers of pesticides and other household chemicals in an age-appropriate manner. Dispose of

unused or expired pesticides properly according to local regulations. By following these recommendations, parents and caregivers can significantly reduce the risk of accidental ingestion of pesticides by children and create a safer home environment. While constant and vigilant supervision is arguably the most effective way to prevent accidental ingestion of harmful substances by children, it is not always feasible in the real world. Parents and caregivers have numerous responsibilities and may not be able to watch their children every second of the day. Children, being naturally curious and explorative, are adept at finding opportunities to investigate their surroundings when adults are momentarily distracted or preoccupied with other tasks. Therefore, relying solely on parental supervision as a preventive measure is not a foolproof solution and can create a false sense of security. It is crucial to acknowledge that even the most attentive parents and caregivers cannot provide uninterrupted supervision at all times. There will inevitably be moments when their attention is diverted, whether it's answering the phone, preparing a meal, or tending to other household chores. During these brief lapses in supervision, children may seize the opportunity to explore cupboards, drawers, or other areas where household chemicals, including pesticides, are stored. Moreover, children's curiosity and determination to explore can sometimes exceed adults' expectations. They may climb onto furniture, access high shelves, or find other ways to reach objects that are seemingly out of reach. This underscores the limitations of relying solely on parental supervision to prevent accidental ingestion. Safe storage practices are therefore essential to minimize the risk of accidental ingestion, even when direct supervision is not possible. By creating a safe environment where hazardous substances are securely stored, parents and caregivers can significantly reduce the likelihood of accidental poisoning. This involves storing pesticides and other household chemicals in locked cabinets or on high shelves, out of reach of children. It also entails using child-resistant packaging and keeping these

substances in their original containers with clear labels and warnings. In addition to safe storage practices, educating children about the potential dangers of pesticides and other household chemicals is crucial. Age-appropriate education can help children understand the risks involved and develop a sense of caution and responsibility. By fostering an understanding of the potential consequences of handling or ingesting these substances, children can learn to make informed decisions and avoid potentially harmful interactions. In essence, while parental supervision is important, it is not a substitute for safe storage practices and education. By combining vigilant supervision with a safe home environment and age-appropriate education, parents and caregivers can create a multi-layered approach to preventing accidental ingestion and protecting children from the harmful effects of pesticides and other household chemicals. Educating parents and caregivers about the potential dangers of pesticides and the importance of safe storage is fundamental in preventing accidental poisoning in children. Public health campaigns, community outreach programs, and educational initiatives can play a vital role in raising awareness and promoting responsible handling and storage of pesticides. By disseminating information about the risks of pesticide exposure and providing guidance on safe storage practices, these programs can empower parents and caregivers to take proactive steps to protect children from accidental poisoning. Parents and caregivers need to understand that pesticides are designed to kill or control pests and can be harmful to humans, especially children. They should be aware of the different types of pesticides and their potential health effects, including acute poisoning and long-term health consequences. Education should emphasize the developmental characteristics of children that make them particularly vulnerable to accidental poisoning. This includes their natural curiosity, tendency to explore their surroundings, and limited understanding of danger and risk. Parents and caregivers should be educated on the importance of safe storage practices to prevent accidental ingestion



of pesticides by children. This includes storing pesticides in their original containers with clear labels, keeping them out of reach of children, and using child-resistant packaging. Education should also focus on recognizing the symptoms of pesticide poisoning in children. Parents and caregivers should be aware of the common signs and symptoms of OP poisoning, such as miosis, hypersalivation, respiratory distress, and seizures. In the event of suspected pesticide poisoning, parents and caregivers should know how to respond promptly and appropriately. This includes calling emergency services, providing basic first aid, and seeking immediate medical attention. Public health campaigns and community outreach programs can effectively disseminate information about the risks of pesticide exposure and promote safe storage practices. These programs can utilize various channels, such as television, radio, print media, and social media, to reach a wide audience. Community health workers and volunteers can also play a crucial role in educating families and communities about pesticide safety. Educational initiatives, such as school programs, workshops, and training sessions, can provide more in-depth information and guidance on pesticide safety. These initiatives can target specific groups, such as parents, caregivers, teachers, and healthcare providers, to enhance their understanding of pesticide risks and prevention strategies. By providing comprehensive education and raising awareness about pesticide safety, these programs can empower parents and caregivers to take proactive steps to protect children from accidental poisoning. They can create a safer home environment, adopt responsible storage practices, and educate their children about the potential dangers of pesticides.<sup>11,12</sup>

The case of the 9-year-old boy who accidentally ingested an organophosphate (OP) insecticide stored in a drinking bottle serves as a poignant example of the vulnerability of children to such incidents. This unfortunate event highlights several critical factors that contribute to accidental poisoning in children and underscores the need for preventive measures to ensure their safety. Children, particularly those under

the age of 10, are at a heightened risk of accidental poisoning due to their developmental stage. Their natural curiosity, coupled with a limited understanding of the hazardous nature of chemicals commonly found in households, makes them particularly susceptible to accidental ingestion. In this case, the child's lack of awareness about the dangerous nature of the substance, coupled with the unsafe storage practice, led to the unfortunate event. The storage of the OP insecticide in a drinking bottle, a seemingly innocuous container, tragically led to the child mistaking it for water, resulting in accidental ingestion. This incident underscores the critical need for safe storage practices for all household chemicals, including pesticides, to prevent accidental poisoning in children. Parents and caregivers must be educated on the importance of keeping potentially harmful substances out of reach of children, in appropriately labeled containers, and ideally under lock and key. This case emphasizes the need for heightened vigilance in protecting children from accidental poisoning. Even with the best intentions, parents and caregivers may not always be able to provide constant supervision. Therefore, it is crucial to create a safe home environment where hazardous substances are securely stored and out of reach of children. Safe storage practices are essential to minimize the risk of accidental ingestion, even when direct supervision is not possible. By creating a safe environment where hazardous substances are securely stored, parents and caregivers can significantly reduce the likelihood of accidental poisoning. This involves storing pesticides and other household chemicals in locked cabinets or on high shelves, out of reach of children. It also entails using child-resistant packaging and keeping these substances in their original containers with clear labels and warnings. Ongoing education is also crucial in preventing accidental poisoning. Parents and caregivers need to be aware of the potential dangers of pesticides and other household chemicals and understand the importance of safe storage practices. Public health campaigns, community outreach programs, and educational

initiatives can play a vital role in raising awareness and promoting responsible handling and storage of pesticides. This case serves as a poignant reminder that accidental poisoning is preventable and that proactive measures can significantly reduce the risk to children. By raising awareness, educating parents and caregivers, and implementing safe storage practices, we can create a safer environment for children and prevent such tragedies from occurring.<sup>13,14</sup>

The successful management of organophosphate (OP) poisoning hinges on the early recognition of its characteristic symptoms and the timely initiation of appropriate medical treatment. These compounds, widely used in agriculture and household pest control, exert their toxicity by inhibiting acetylcholinesterase, an enzyme crucial for the breakdown of the neurotransmitter acetylcholine. This inhibition leads to an accumulation of acetylcholine at nerve synapses, causing a wide range of symptoms due to overstimulation of both muscarinic and nicotinic receptors. Muscarinic symptoms are prominent in OP poisoning and include miosis (constriction of the pupils), hypersalivation, hyperlacrimation (excessive tearing), urination, defecation, gastrointestinal distress (nausea, vomiting, and abdominal cramps), emesis (vomiting), bronchorrhea (excessive bronchial secretions), bronchospasm (constriction of the airways), and bradycardia (slow heart rate). Nicotinic symptoms, on the other hand, include muscle fasciculations (twitching), muscle weakness, tachycardia (rapid heart rate), hypertension (high blood pressure), and paralysis. Early recognition of these symptoms is paramount in initiating prompt and effective treatment. In the case of the 9-year-old boy presented earlier, the presence of classic muscarinic symptoms such as miosis, hypersalivation, and respiratory distress, along with the reported history of ingestion, allowed the medical team to rapidly diagnose OP poisoning and commence treatment without delay. Atropine, a muscarinic receptor antagonist, is the cornerstone of treatment for the muscarinic symptoms of OP poisoning. It acts

by competitively blocking the binding of acetylcholine to muscarinic receptors, effectively counteracting the excessive cholinergic stimulation and alleviating the associated symptoms. The prompt administration of atropine in this case led to a significant improvement in the child's condition, highlighting the efficacy of this antidote in reversing the muscarinic effects of OP poisoning. However, it is crucial to emphasize that atropine dosage must be carefully titrated to effect, as inadequate atropinization can lead to treatment failure and potentially life-threatening complications. The initial dose of atropine should be administered intravenously and repeated every 5-10 minutes until the muscarinic symptoms, such as excessive secretions and bronchospasm, subside. The patient's heart rate and pupil size should be closely monitored to assess the adequacy of atropinization. In addition to atropine, supportive care plays a vital role in the management of OP poisoning. This includes ensuring adequate oxygenation through oxygen therapy, maintaining hydration and electrolyte balance with intravenous fluids, and providing respiratory support with mechanical ventilation if necessary. Continuous monitoring of vital signs, oxygen saturation, and urine output is essential to assess the patient's response to treatment and detect any complications promptly. Early recognition and prompt management of OP poisoning are crucial to prevent severe complications and improve patient outcomes. The timely administration of atropine, along with appropriate supportive care, can effectively reverse the muscarinic effects of OP poisoning and prevent life-threatening complications such as respiratory failure and cardiovascular collapse.<sup>15,16</sup>

In addition to the administration of atropine, supportive care plays a crucial role in the comprehensive management of organophosphate (OP) poisoning. Supportive care aims to stabilize the patient's physiological functions, prevent complications, and provide an optimal environment for recovery. In the case of the 9-year-old boy, the supportive care provided in the pediatric intensive care unit (PICU), along with the timely administration

of atropine, contributed significantly to the complete resolution of his symptoms and his eventual recovery. OP poisoning can cause respiratory complications due to the accumulation of secretions in the airways, bronchospasm, and weakness of respiratory muscles. Supportive care may include oxygen therapy to maintain adequate oxygenation, suctioning to clear airway secretions, and mechanical ventilation if respiratory failure ensues. OP poisoning can also affect the cardiovascular system, leading to bradycardia (slow heart rate), hypotension (low blood pressure), and even cardiac arrest in severe cases. Supportive care may involve monitoring heart rate and blood pressure, administering intravenous fluids to maintain blood volume and circulation, and providing medications to support heart function if necessary. Vomiting and diarrhea, common symptoms of OP poisoning, can lead to dehydration and electrolyte imbalances. Supportive care includes administering intravenous fluids to maintain hydration and electrolyte balance, correcting any imbalances, and monitoring urine output to assess kidney function. If the OP pesticide was ingested, gastrointestinal decontamination may be necessary to remove any remaining toxin from the stomach. This may involve gastric lavage (washing out the stomach) or administering activated charcoal to bind the toxin and prevent its absorption. OP poisoning can affect the patient's ability to eat and digest food. Supportive care may include providing nutritional support through intravenous fluids or enteral feeding (feeding through a tube into the stomach or intestines) to ensure adequate nutrition during recovery. Continuous monitoring of vital signs, oxygen saturation, and urine output is essential to assess the patient's response to treatment and detect any complications promptly. Regular neurological assessments are also crucial to monitor for any signs of seizures or other neurological complications. Supportive care plays a vital role in the management of OP poisoning. Maintaining physiological stability and preventing complications. Providing an optimal environment for recovery. Reducing the severity and duration of symptoms.

Improving patient outcomes. In the case of the 9-year-old boy, the supportive care provided in the PICU, including oxygen therapy, intravenous fluids, and continuous monitoring, contributed significantly to his recovery. The close monitoring and prompt management of any emerging complications ensured that his condition remained stable and allowed for the complete resolution of his symptoms.<sup>17,18</sup>

The case of the 9-year-old boy who accidentally ingested an OP insecticide serves as a poignant reminder of the urgent need for comprehensive public health interventions aimed at preventing such incidents. These interventions should focus on multiple levels, including education and awareness, policy and regulation, and community engagement, to effectively reduce the risk of accidental OP poisoning in children. Educating parents, caregivers, and the general public about the potential dangers of pesticides and the importance of safe storage and handling practices is paramount. Public health campaigns should utilize various channels, such as television, radio, print media, and social media, to disseminate information about the risks of pesticide exposure and provide clear guidance on safe storage and handling practices. Community health workers and volunteers can also play a crucial role in educating families and communities about pesticide safety. Emphasize that pesticides are designed to be toxic to living organisms and can pose serious health risks to humans, especially children. Highlight the developmental characteristics of children that make them more susceptible to accidental poisoning, such as their curiosity, exploratory behavior, and limited understanding of danger. Provide clear and concise instructions on safe storage practices, including keeping pesticides in their original containers with labels intact, storing them out of reach of children in locked cabinets or on high shelves, and using child-resistant packaging. Educate parents and caregivers about the common signs and symptoms of OP poisoning, enabling them to identify potential poisoning early and seek prompt medical attention. Provide guidance on basic first aid measures and

emergency response in case of suspected pesticide poisoning. Effective policies and regulations are essential to control the availability and accessibility of highly hazardous pesticides. Governments should implement stricter regulations on the sale and distribution of OP pesticides, limiting their availability for household use and promoting safer alternatives. Policies should also mandate child-resistant packaging and clear labeling with appropriate warnings and instructions for use. Engaging communities in pesticide safety efforts is crucial for creating a supportive environment that promotes safe practices. Community-based programs can organize workshops, training sessions, and awareness campaigns to educate residents about pesticide risks and prevention strategies. Local leaders and community organizations can play a vital role in advocating for safer pesticide practices and promoting community-wide initiatives to reduce the risk of accidental poisoning. Whenever feasible, promoting the use of safer alternatives to OP insecticides can further contribute to minimizing the risk of poisoning. Integrated pest management strategies, which emphasize non-chemical methods of pest control, can reduce reliance on pesticides. Encouraging the use of less toxic pesticides, such as biopesticides or those with lower toxicity profiles, can also help minimize the risk of severe poisoning.<sup>19,20</sup>

#### 4. Conclusion

This case report underscores the critical importance of early recognition and prompt management of OP poisoning in children. The timely administration of atropine, along with appropriate supportive care, can effectively reverse the muscarinic effects of OP poisoning and prevent life-threatening complications. This case also serves as a stark reminder of the urgent need for public health interventions aimed at preventing accidental OP poisoning in children. Educating parents and caregivers about the safe storage and handling of OP insecticides is paramount in reducing the risk of accidental ingestion. By raising awareness and

implementing preventive measures, we can strive to reduce the incidence of this preventable tragedy and safeguard children from the harmful effects of OP pesticides.

#### 5. References

1. Lau CL, Chung KL, Kam CW. Accidental organophosphate insecticide poisoning: a fatal child case with hazards to emergency department staff. *Hong Kong J Emerg Med.* 2003; 10(4): 248–52.
2. Wheeler DS, Fisher LE, Mehta R, Segeleon JE, Pearson-Shaver AL. Organophosphate poisoning in children. *Crit Care Med.* 2004; 32(Suppl): A180.
3. Kiat W. Acute deliberate organophosphate (Coumaphos) poisoning with intermediate syndrome in a 1 year old child. *Toxicol Lett.* 2007; 172: S132–3.
4. El-Naggar AE-R, Abdalla MS, El-Sebaey AS, Badawy SM. Clinical findings and cholinesterase levels in children of organophosphates and carbamates poisoning. *Eur J Pediatr.* 2009; 168(8): 951–6.
5. Pavlovic M, Neubauer D, Al-Tawari AA. Delayed effects of transcutaneous organophosphate poisoning in four children. *Child Neurol Open.* 2015; 2(4): 2329048X15618970.
6. Koirala DP, Rao KS, Malla KK, Malla T. A study of clinical features, management and outcome of organophosphate and Carbamate poisoning in children. *J Nepal Paediatr Soc.* 2013; 33(2): 85–90.
7. Donald KA, Adnams CM, Rother A, Tsze D, Brand D, London L. P102 – 2305: Long- term neurodevelopmental and behavioural outcomes of acute organophosphate poisoning in South African children. *Eur J Paediatr Neurol.* 2015; 19: S123.
8. Hamad MH, Adeel AA, Alhaboob AAN, Ashri AM, Salih MA. Acute poisoning in a child following topical treatment of head lice

- (pediculosis capitis) with an organophosphate pesticide. *Sudan J Paediatr.* 2016; 16(1): 63–6.
9. Yesilbas O, Kihdir HS, Altiti M, Petmezci MT, Balkaya S, Bursal Duramaz B, et al. Acute severe organophosphate poisoning in a child who was successfully treated with therapeutic plasma exchange, high-volume hemodiafiltration, and lipid infusion. *J Clin Apher.* 2016; 31(5): 467–9.
  10. Ijezie E, Megbelayin F, Edem K, Ijezie AE. Accidental organophosphate poisoning in a child in Uyo, Nigeria: a public health alert. *Int J Sci Rep.* 2016; 2(5): 106.
  11. Balme K, McCulloch M, Stephen C. Prolonged paralysis in a child with organophosphate pesticide poisoning. *S Afr Med J.* 2018; 108(6): 468–70.
  12. Dhanalakshmi, Febina, Sabapathy S, Chaithra, Thilepan. Correlation of serum creatine phosphokinase and serum cholinesterase in organophosphate poisoning in children. *Int J Contemp Pediatr.* 2019; 6(6): 2352.
  13. Sharma N, Nin-Gonzalez R. Organophosphate poisoning in a young child: a case report. *Oxf Med Case Reports.* 2021; 2021(2): omaa137.
  14. Abdel Baseer KA, Gad EF, Abdel Raheem YF. Clinical profile and outcome of acute organophosphate poisoning in children of Upper Egypt: a cross-sectional study. *BMC Pediatr.* 2021; 21(1): 98.
  15. Joson MVASG, Castor FRM, Micu-Oblefias CV. Role of intravenous lipid emulsion therapy and packed red blood cell transfusion as adjuvant treatment in the management of a child with severe organophosphate poisoning (chlorpyrifos). *BMJ Case Rep.* 2022; 15(4): e246381.
  16. Sailan A, Lekshmi A, Babu B. Organophosphate poisoning in a young child- A case report. *Asian J Pediatr Res.* 2022; 18–20.
  17. Manu Prakash S K, Kumar S R, Harish H N, Suman S. Clinico-epidemiological profile of acute pancreatitis secondary to organophosphate poisoning in children at a tertiary care centre: Case series. *Indian J Public Health Res Dev.* 2023; 14(4): 126–31.
  18. Althaqafi W, Alanazi RI, Almeshari WK. The use of amantadine in treating extrapyramidal symptoms in organophosphates poisoning in a child. *Case Rep Pediatr.* 2023; 2023: 1632052.
  19. Patil MM, Doddihal CR, Sinha A, Kumar Bm P. Unwashed grapes as a cause of organophosphate poisoning in a child in rural India. *BMJ Case Rep.* 2024; 17(8): e260702.
  20. Mwaba C, Chungu C, Chola R, Nkole KL, Wa Somwe S, Mpabalwani E. Organophosphate insecticide poisoning with monocrotophos-induced fabricated illness in a 7-year-old girl with refractory seizures over a 4-year period. *Paediatr Int Child Health.* 2022; 42(2): 83–8.