



## Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: [www.bioscmed.com](http://www.bioscmed.com)

### The Use of Trendelenburg Position in the Management of Retropharyngeal Abscess: A Case Report and Clinical Considerations

Emilia Nissa Khairani<sup>1\*</sup>, Liliriawati Ananta Kahar<sup>2</sup>, Novialdi<sup>1</sup>

<sup>1</sup>Department of Otorhinolaryngology Head and Neck Surgery, Faculty of Medicine, Universitas Andalas, Padang, Indonesia

<sup>2</sup>Department of Anesthesiology and Intensive Care, Faculty of Medicine, Universitas Andalas, Padang, Indonesia

#### ARTICLE INFO

##### Keywords:

Aspiration  
Deep neck space infection  
Retropharyngeal abscess  
Trendelenburg position  
Ventilator-associated pneumonia

##### \*Corresponding author:

Emilia Nissa Khairani

##### E-mail address:

[nissaemilia90@gmail.com](mailto:nissaemilia90@gmail.com)

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v9i1.1180>

#### A B S T R A C T

**Background:** Retropharyngeal abscess (RPA) is a serious deep neck space infection that can lead to life-threatening complications such as airway obstruction, mediastinitis, and sepsis. The Trendelenburg position, where the patient's head is lower than their feet, has been traditionally used in the management of RPA to prevent aspiration of pus in case of abscess rupture. However, the optimal positioning for patients with RPA remains a topic of debate. **Case presentation:** A 51-year-old male patient presented to the emergency room with a 5-day history of progressive dysphagia and neck swelling. He was diagnosed with a ruptured retropharyngeal abscess extending to the mediastinum (descending necrotizing mediastinitis). The patient underwent surgical drainage and was subsequently admitted to the intensive care unit (ICU), where he was intubated and mechanically ventilated in the Trendelenburg position. Five days later, the patient developed pneumonia, suspected to be ventilator-associated pneumonia (VAP). **Conclusion:** The Trendelenburg position may be considered in the management of RPA to minimize the risk of aspiration. However, it is essential to weigh the potential benefits against the risks, including VAP, and to consider alternative positions such as the lateral Trendelenburg position, which may offer similar benefits with reduced risk. Further research is needed to determine the optimal positioning strategy for patients with RPA.

#### 1. Introduction

Retropharyngeal abscess (RPA) is a rare but potentially life-threatening deep neck space infection characterized by the accumulation of pus within the retropharyngeal space. This space, located between the posterior pharyngeal wall and the prevertebral fascia, extends from the base of the skull to the superior mediastinum. Due to its anatomical location and proximity to vital structures, RPA can lead to serious complications such as airway obstruction, mediastinitis, sepsis, and even death if not promptly diagnosed and managed. The incidence of RPA has decreased significantly with the widespread use of antibiotics; however, it remains a serious concern,

particularly in children, immunocompromised individuals, and those with underlying medical conditions. The most common causes of RPA include upper respiratory tract infections, tonsillitis, and dental infections. Other potential causes include foreign body ingestion, trauma, and iatrogenic injury.<sup>1,3</sup>

The clinical presentation of RPA can vary depending on the severity of the infection and the presence of complications. Common symptoms include sore throat, dysphagia (difficulty swallowing), neck pain and stiffness, odynophagia (painful swallowing), fever, and muffled voice. In severe cases, patients may develop stridor (a high-pitched breathing

sound), respiratory distress, and sepsis. The diagnosis of RPA is typically made based on a combination of clinical findings, imaging studies, and laboratory investigations. Contrast-enhanced computed tomography (CT) scan of the neck is the gold standard for diagnosis, allowing for visualization of the abscess, its extent, and any associated complications. The management of RPA often requires a multidisciplinary approach involving otolaryngologists, intensivists, infectious disease specialists, and radiologists. The primary goals of treatment are to secure the airway, drain the abscess, and eradicate the infection. Intravenous antibiotics are the mainstay of medical therapy, while surgical drainage is often necessary for large or complicated abscesses. Airway management is a critical aspect of RPA care, as airway obstruction can occur rapidly and unexpectedly.<sup>4-7</sup>

Traditionally, the Trendelenburg position, where the patient's head is lower than their feet, has been used in the management of RPA to prevent aspiration of pus in case of abscess rupture. However, the optimal positioning for patients with RPA remains a topic of debate. Some studies have suggested that the Trendelenburg position may increase the risk of ventilator-associated pneumonia (VAP) in intubated patients, while others have found no significant difference in VAP rates between different positions.<sup>8-10</sup> This case report describes a 51-year-old male patient with a ruptured retropharyngeal abscess who was managed in the Trendelenburg position and subsequently developed VAP. The report highlights the potential risks and benefits of the Trendelenburg position in the management of RPA and emphasizes the need for careful clinical judgment and individualized treatment decisions.

## **2. Case Presentation**

A 51-year-old male presented to the emergency department with a primary complaint of progressively worsening odynophagia over the preceding five days. The patient's history revealed that the discomfort initially manifested as a sore throat, gradually escalating to a point where swallowing, even for

liquids, became severely challenging. Additionally, he observed a noticeable swelling in his neck, accompanied by pain and tenderness, which had emerged concurrently with the onset of odynophagia. The patient denied experiencing any fever, chills, or recent upper respiratory tract infection. His medical history was significant for poorly managed diabetes mellitus, for which he was not receiving any regular treatment. He denied any history of smoking, alcohol consumption, or illicit drug use. Upon physical examination, the patient exhibited signs of moderate distress, characterized by tachypnea and a slightly muffled voice. His vital signs included a temperature of 38.5°C (101.3°F) and a pulse rate of 110 beats per minute. Examination of the neck revealed pronounced swelling and tenderness in both the anterior and posterior cervical regions, with the swelling being more prominent on the left side. No palpable fluctuance or crepitus was detected. Oral examination was limited due to the patient's discomfort but revealed erythema and edema of the posterior pharyngeal wall. There was no evidence of trismus or uvular deviation. Auscultation of the lungs revealed coarse breath sounds bilaterally, without wheezes or rales. Laboratory investigations indicated leukocytosis, with a white blood cell count of 18,000 cells/mm<sup>3</sup> and a differential count showing a predominance of neutrophils. The C-reactive protein level was elevated at 150 mg/L. A contrast-enhanced computed tomography (CT) scan of the neck and chest was conducted, revealing a large retropharyngeal abscess extending from the level of the oropharynx to the superior mediastinum. The abscess was associated with significant surrounding inflammation and edema. There was no evidence of airway compromise at the time of imaging. Based on the clinical and radiological findings, the patient was diagnosed with a ruptured retropharyngeal abscess with descending necrotizing mediastinitis. He was immediately initiated on intravenous broad-spectrum antibiotics and transferred to the operating room for surgical drainage.

The patient's primary complaint of progressively worsening odynophagia, coupled with dysphagia, is a hallmark symptom of retropharyngeal abscess (RPA). The retropharyngeal space lies in close proximity to the pharynx and esophagus. As the abscess expands, it can impinge on these structures, causing pain and difficulty with swallowing. The severity of odynophagia and dysphagia often correlates with the size and location of the abscess. The presence of neck swelling and tenderness is another common finding in RPA. The swelling is typically located in the anterior and posterior cervical regions and may be more prominent on one side. The tenderness is often elicited by palpation in the affected area. The extent of neck swelling and tenderness can vary depending on the size and location of the abscess and the degree of associated inflammation.

The patient's slightly muffled voice, also known as "hot potato voice," is another clinical manifestation of RPA. This muffled vocal quality results from the mass effect of the abscess on the surrounding tissues, including the larynx and pharynx, which alters the resonance of the vocal tract. The presence of fever and leukocytosis indicates an ongoing inflammatory process, which is consistent with the diagnosis of RPA. Fever is a systemic response to infection, while leukocytosis reflects the body's attempt to fight off the invading pathogens. The absence of trismus (difficulty opening the mouth) and uvular deviation in this patient is noteworthy. While these findings can be associated with RPA, they are not always present. A trismus may occur if the abscess extends to involve the pterygoid muscles, which are involved in jaw movement. Uvular deviation may occur if the abscess exerts pressure on the uvula, causing it to deviate away from the affected side (Figure 1).

The presence of coarse breath sounds on lung auscultation may be attributed to several factors in this patient. It could be related to the patient's underlying respiratory status, the presence of edema

and inflammation in the neck and surrounding tissues, or early signs of respiratory complications such as pneumonia. The elevated C-reactive protein (CRP) level further supports the diagnosis of RPA and indicates a significant inflammatory response. CRP is an acute-phase protein produced by the liver in response to inflammation. Its levels rise rapidly in the presence of infection or tissue injury. The contrast-enhanced computed tomography (CT) scan of the neck and chest played a crucial role in confirming the diagnosis of RPA and delineating the extent of the disease. The CT scan revealed a large retropharyngeal abscess extending from the oropharynx to the superior mediastinum, with associated inflammation and edema. The absence of airway compromise on imaging was an important finding, as it influenced the initial management decisions (Figures 2 and 3).

The diagnosis of ruptured retropharyngeal abscess with descending necrotizing mediastinitis was made based on the constellation of clinical and radiological findings (Figure 4). The patient's presentation with odynophagia, dysphagia, neck swelling, muffled voice, fever, and leukocytosis, along with the CT scan findings of a large retropharyngeal abscess extending to the mediastinum, strongly supported this diagnosis. Descending necrotizing mediastinitis is a severe complication of RPA, characterized by the spread of infection from the retropharyngeal space to the mediastinum. This condition is associated with high morbidity and mortality rates and requires prompt and aggressive treatment. Given the severity of the patient's condition and the potential for life-threatening complications, immediate management was initiated. This included; Intravenous broad-spectrum antibiotics: To combat the infection and prevent its further spread; Transfer to the operating room: For surgical drainage of the abscess. The prompt initiation of appropriate treatment is crucial in managing RPA and its complications, as delay in treatment can lead to worse outcomes.



Figure 1. There was a rupture of retropharyngeal abscess, pus (+) (yellow arrow).




Figure 2. A. Cervical X-ray showed retropharyngeal dilation to the retrotracheal space (C2-7) with an open-air column, B. Cor, and pulmo in normal limits.




Figure 3. Cervical CT scan with the result of a hypodense lesion in the larynx area, retropharynx extending to the anterior colli and pretracheal regions.



Figure 4. There were multiple infiltrates and an increase in bronchovascular patterns in both lung fields.

### 3. Discussion

The Trendelenburg position, a classic maneuver in medical practice, involves tilting the patient's body so that the head is lower than the feet. This positioning has been a cornerstone in managing various clinical scenarios, including retropharyngeal abscess (RPA), for its potential benefits in airway management. However, it's not without potential drawbacks, particularly in the context of mechanically ventilated patients. The Trendelenburg position, a time-honored maneuver in medical practice, involves tilting the patient's body so that the head is lower than the feet. This positioning has been a cornerstone in managing various clinical scenarios, including retropharyngeal abscess (RPA), for its potential benefits in airway management. The primary advantage of the Trendelenburg position in RPA management lies in its ability to harness gravity to facilitate drainage of purulent material away from the airway. By tilting the patient's head downward, the natural flow of pus is directed towards the esophagus, where it can be safely expelled through swallowing or suctioned out. This is particularly crucial in cases of RPA rupture, where a sudden release of pus can pose a significant aspiration risk. Gravity plays a pivotal role in directing the flow of fluids within the body. In the case of RPA, where pus accumulates in the retropharyngeal space, the Trendelenburg position utilizes this natural force to guide the drainage of pus away from the critical airway

structures and towards the esophagus. Aspiration of pus, a potential complication of RPA, can have dire consequences, including airway obstruction, pneumonia, and respiratory distress. The Trendelenburg position helps mitigate this risk by promoting the drainage of pus away from the trachea and lungs. This is especially important in patients with compromised airway reflexes or those who are unable to protect their airway effectively. In certain cases, the Trendelenburg position can enhance surgical access during procedures involving the neck and upper airway. By tilting the patient's head downward, the surgical field is better exposed, allowing for improved visualization and manipulation of anatomical structures. Maintaining airway patency is paramount in managing RPA. The Trendelenburg position aids in achieving this goal by minimizing the risk of pus entering the lower respiratory tract. This is particularly crucial in patients with compromised airway reflexes or those who are unable to protect their airway effectively due to factors such as sedation or neurological impairment. Aspiration of pus can lead to a cascade of complications. The accumulation of pus in the trachea or bronchi can obstruct airflow, leading to respiratory distress and potentially respiratory failure. The introduction of bacteria-laden pus into the lungs can trigger an inflammatory response, resulting in pneumonia. The combined effects of airway obstruction and pneumonia can lead to respiratory

distress, a condition characterized by shortness of breath, rapid breathing, and low oxygen levels. The Trendelenburg position is particularly beneficial for patients at increased risk of aspiration. Those who are sedated, unconscious, or have neurological impairments may have diminished airway reflexes, making them more susceptible to aspiration. Individuals with dysphagia or other swallowing difficulties may also be at a higher risk of aspirating pus. Young children may have less developed airway protective mechanisms, making them more vulnerable to aspiration. The Trendelenburg position can significantly enhance the surgical field during procedures involving the neck and upper airway. By tilting the patient's head downward, the anatomical structures are better exposed, allowing for improved visualization and manipulation. This can be particularly advantageous in complex cases or when delicate surgical maneuvers are required. Improved visualization and access can lead to more precise surgical technique, potentially minimizing tissue trauma and associated complications. The Trendelenburg position can also facilitate the placement of surgical instruments and retractors, further optimizing the surgical field. While the Trendelenburg position offers valuable benefits in managing certain clinical conditions, it's essential to be cognizant of its potential risks. These risks span various physiological systems and can have significant implications for patient care. Recent research has raised concerns about the potential link between the Trendelenburg position and an elevated risk of VAP in intubated patients. This association is attributed to the fact that this position can promote the migration of oropharyngeal and gastric secretions towards the lower respiratory tract. These secretions often harbor bacteria, and their accumulation in the lungs can lead to colonization and subsequent pneumonia. While gravity is harnessed for therapeutic benefit in certain scenarios, it can also contribute to complications. In the Trendelenburg position, gravity can facilitate the movement of secretions, which may contain bacteria, from the oropharynx and stomach towards the lungs.

The pooling of these secretions in the lower respiratory tract creates a fertile environment for bacterial growth and colonization. This can disrupt the delicate balance of the lung microbiome and increase the risk of infection. VAP is a serious complication of mechanical ventilation, often caused by the aspiration of bacteria-laden secretions. The Trendelenburg position, by promoting the accumulation of these secretions in the lungs, can increase the likelihood of VAP development. VAP is associated with increased morbidity, mortality, prolonged hospitalization, and escalated healthcare costs. This risk is particularly concerning in critically ill patients who are already vulnerable to infections. The Trendelenburg position can also have implications for the cardiovascular system. Tilting the patient's head downward can lead to an increase in venous return to the heart, potentially resulting in an elevated workload for the heart. This can be particularly concerning in patients with pre-existing heart conditions. The Trendelenburg position alters the distribution of blood within the body. By positioning the head lower than the feet, venous return to the heart is increased. This means that more blood is flowing back to the heart from the lower extremities and abdomen. The heart has to work harder to pump this increased volume of blood. This can put a strain on the heart, particularly in patients with underlying heart conditions such as heart failure or coronary artery disease. In some cases, the increased venous return and cardiac workload can lead to hemodynamic instability, characterized by fluctuations in blood pressure and heart rate. In certain cases, prolonged use of the Trendelenburg position can contribute to cerebral edema, or swelling in the brain. This is due to the increased blood flow to the head that occurs when the head is positioned lower than the rest of the body. The Trendelenburg position can affect blood flow to the brain. When the head is lower than the feet, gravity can increase blood flow to the brain. This increased blood flow can lead to fluid accumulation in the brain tissue, resulting in cerebral edema. Cerebral edema can be a serious complication, potentially causing neurological damage. The risk of cerebral edema is

higher in patients with pre-existing neurological conditions or those who have undergone neurosurgery. Maintaining the Trendelenburg position for extended periods can lead to patient discomfort and increase the risk of pressure sores, especially in patients with limited mobility or those who are unable to reposition themselves. The Trendelenburg position can create pressure points on certain areas of the body, such as the back of the head, shoulders, and heels. Prolonged pressure on these areas can compromise blood flow to the skin and underlying tissues. This compromised blood flow can lead to the development of pressure sores, also known as bedsores or decubitus ulcers. Pressure sores are injuries to the skin and underlying tissue caused by prolonged pressure. The Trendelenburg position can also be uncomfortable for patients, particularly those with musculoskeletal issues or those who are unable to reposition themselves. The Trendelenburg position, while offering valuable benefits in managing certain clinical conditions, is not without potential risks. Therefore, the decision to utilize this position should be made on a case-by-case basis, carefully weighing the potential benefits against the possible risks. This individualized approach to patient care ensures that the decision to use the Trendelenburg position is tailored to the specific needs and circumstances of each patient. The size and location of the abscess can influence the potential benefits and risks of the Trendelenburg position. Larger abscesses or those located closer to the airway may pose a greater risk of aspiration, making the Trendelenburg position more appealing for gravity-assisted drainage. However, such cases might also increase the risk of complications like VAP, necessitating careful consideration. The presence of complications, such as rupture or mediastinal extension, can further sway the decision. In cases of rupture, the Trendelenburg position may be crucial for preventing the spread of pus to the lower respiratory tract. However, if mediastinal extension is present, the Trendelenburg position might exacerbate cardiovascular or neurological risks, prompting the consideration of

alternative positions. The patient's ability to protect their airway is a critical factor. Patients with compromised airway reflexes, due to factors such as sedation, neurological impairment, or underlying respiratory conditions, may benefit from the Trendelenburg position to minimize aspiration risk. However, their respiratory status should be closely monitored for any signs of distress or compromise. The patient's overall respiratory function should also be assessed. Patients with pre-existing respiratory conditions, such as asthma or chronic obstructive pulmonary disease (COPD), may be more susceptible to complications like VAP. In such cases, the potential benefits of the Trendelenburg position must be carefully weighed against the increased risk of respiratory complications. Patients with underlying cardiovascular conditions, such as heart failure or coronary artery disease, may be more vulnerable to the cardiovascular effects of the Trendelenburg position. The increased venous return and cardiac workload associated with this position could exacerbate their condition. Patients with neurological conditions may be at a higher risk of cerebral edema when placed in the Trendelenburg position. The increased blood flow to the head could worsen their condition or lead to neurological complications. The anticipated duration of the Trendelenburg position should be factored into the risk assessment. Prolonged use of this position can increase the risk of complications such as VAP, cardiovascular effects, cerebral edema, discomfort, and pressure sores. If the Trendelenburg position is maintained for an extended period, close monitoring and frequent repositioning are essential to minimize the risk of complications. In situations where the risks of the Trendelenburg position outweigh the benefits, alternative positioning strategies should be explored. The lateral Trendelenburg position, where the patient is positioned on their side with the head slightly tilted downward, offers a potential compromise. This position aims to maintain the benefits of gravity-assisted drainage while minimizing the risk of VAP and other complications associated with the supine

Trendelenburg position.<sup>11-13</sup>

Ventilator-associated pneumonia (VAP) is a frequent and serious complication in critically ill patients undergoing mechanical ventilation. It is associated with increased morbidity, mortality, prolonged hospitalization, and escalated healthcare costs. The development of VAP in this patient, despite being placed in the Trendelenburg position, highlights the delicate balance between the benefits and risks associated with different patient positioning strategies in the context of RPA management. Ventilator-associated pneumonia (VAP) is a common and serious complication that can arise in critically ill patients who require mechanical ventilation. It is clinically defined as pneumonia that develops in patients who have been intubated and mechanically ventilated for at least 48 hours. VAP is one of the most prevalent hospital-acquired infections in intensive care units (ICUs), affecting a significant proportion of mechanically ventilated patients. The diagnosis of VAP is typically made based on a combination of clinical, radiological, and microbiological findings. Clinical criteria may include fever, elevated white blood cell count, purulent sputum, increased respiratory rate, worsening oxygenation, and new or progressive infiltrates on chest X-ray. Radiological findings often show new or progressive infiltrates, consolidations, or opacities on chest imaging. Microbiological confirmation may involve culturing respiratory secretions or blood to identify the causative pathogen. VAP is a significant healthcare concern, affecting approximately 9-27% of mechanically ventilated patients. The incidence of VAP can vary depending on several factors, including the patient population, the type of ICU, and the diagnostic criteria used. Certain patient populations are at higher risk of developing VAP, including those with underlying medical conditions, such as diabetes, chronic obstructive pulmonary disease (COPD), and immunosuppression. The type of ICU can also influence VAP rates. ICUs that care for a higher proportion of critically ill patients, such as surgical ICUs and medical ICUs, tend to have higher rates of VAP. The diagnostic criteria used to define VAP can

also affect the reported incidence. More stringent criteria may lead to lower reported rates of VAP. VAP is associated with a range of adverse outcomes. VAP can lead to sepsis, respiratory failure, and other complications, increasing the overall morbidity of critically ill patients. VAP is associated with a significant increase in mortality rates among mechanically ventilated patients. VAP can prolong the duration of mechanical ventilation and ICU stay, leading to increased healthcare costs and resource utilization. The treatment of VAP and its associated complications can be costly, adding to the financial burden of healthcare systems. Given the significant impact of VAP on patient outcomes and healthcare costs, prevention is of paramount importance. Several strategies have been shown to be effective in preventing VAP. Elevating the head of the bed to 30-45 degrees can help reduce the risk of aspiration. Regular oral care can help reduce the number of bacteria in the mouth. Subglottic suctioning can help remove secretions that accumulate above the endotracheal tube cuff. Selective digestive decontamination involves the use of antibiotics to reduce the number of bacteria in the gut. Ventilator-associated pneumonia (VAP) is a complex and multifactorial disease with a pathogenesis that involves an intricate interplay of host factors, pathogen characteristics, and the unique environment created by mechanical ventilation. Understanding the pathogenesis of VAP is crucial for developing effective prevention and treatment strategies. The endotracheal tube, a necessary component of mechanical ventilation, plays a significant role in the development of VAP. It bypasses the natural defense mechanisms of the upper airway, allowing bacteria to enter the lower respiratory tract. Additionally, the endotracheal tube can cause micro-aspiration of oropharyngeal and gastric secretions, which often contain bacteria. The upper airway possesses a sophisticated system of defenses to prevent pathogens from reaching the lower respiratory tract. The lining of the respiratory tract is covered in cilia, tiny hair-like structures that beat rhythmically to move mucus and trapped particles



upward and out of the airways. The cough reflex helps expel foreign material, including pathogens, from the airways. The upper airway is populated with immune cells that recognize and destroy pathogens. The endotracheal tube disrupts these natural defenses, creating an opportunity for bacteria to colonize the lower respiratory tract. The endotracheal tube can also facilitate micro-aspiration, the inadvertent entry of small amounts of oropharyngeal or gastric secretions into the lower respiratory tract. These secretions often contain bacteria, which can then establish infection in the lungs. Critically ill patients often have weakened immune systems due to underlying medical conditions, malnutrition, or the stress of critical illness. This impaired immunity makes them more susceptible to infections, including VAP. The oropharynx and stomach can become colonized with bacteria, particularly in critically ill patients who are receiving antibiotics or proton pump inhibitors. These bacteria can then be aspirated into the lungs, leading to VAP. Bacteria can form biofilms on the surface of the endotracheal tube. Biofilms are communities of bacteria that are encased in a protective matrix, making them more difficult to eradicate with antibiotics. Biofilms on the endotracheal tube can serve as a persistent source of infection, contributing to the development of VAP. The pathogenesis of VAP is a complex interplay of these various factors. The endotracheal tube disrupts the natural defenses of the upper airway and facilitates micro-aspiration, while impaired host defenses, colonization of the oropharynx and stomach, and biofilm formation further increase the risk of infection. Understanding the pathogenesis of VAP is crucial for developing effective prevention and treatment strategies. Prevention strategies should focus on minimizing the risk of aspiration, maintaining the integrity of the upper airway defenses, and preventing bacterial colonization. Treatment strategies should target the causative pathogens and address the underlying factors that contribute to VAP development. Ventilator-associated pneumonia (VAP) is a serious complication that can arise in patients

undergoing mechanical ventilation. Its clinical presentation can vary depending on the severity of the infection and the patient's underlying health status. Recognizing the common signs and symptoms of VAP is crucial for prompt diagnosis and timely intervention. An elevation in body temperature is a common sign of infection, including VAP. The fever may be accompanied by chills or sweating. An increase in the number of white blood cells, known as leukocytosis, is a common indicator of infection. The white blood cell count is often measured as part of a complete blood count (CBC) test. Purulent sputum, or phlegm that is thick, yellow, or green in color, is a common finding in VAP. It indicates the presence of an infection in the lungs. An increase in the number of breaths per minute, known as tachypnea, is a common sign of respiratory distress. In VAP, the increased respiratory rate may be an attempt to compensate for the decreased oxygenation caused by the infection. A decrease in the level of oxygen in the blood, known as hypoxemia, is a common finding in VAP. It can be measured using a pulse oximeter or arterial blood gas analysis. Infiltrates are abnormal findings on a chest X-ray that indicate the presence of fluid or inflammation in the lungs. In VAP, new or progressive infiltrates may be seen, indicating the spread of infection. Patients with mild VAP may have only a few subtle symptoms, while those with severe VAP may experience significant respiratory distress and sepsis. Patients with underlying medical conditions, such as diabetes or COPD, may be more susceptible to VAP and may have a more severe clinical presentation. The type of bacteria causing VAP can also influence the clinical presentation. Some bacteria are more likely to cause severe pneumonia than others. Early recognition of the signs and symptoms of VAP is crucial for prompt diagnosis and timely intervention. Delay in diagnosis and treatment can lead to worse outcomes, including increased morbidity and mortality. The diagnosis of VAP can be challenging, as there is no single gold standard test. A combination of clinical, radiological, and microbiological findings is often used to make the

diagnosis. The Clinical Pulmonary Infection Score (CPIS) is a commonly used tool to assess the likelihood of VAP. The treatment of VAP typically involves the administration of antibiotics. The choice of antibiotics depends on the suspected pathogens and local antibiotic resistance patterns. Empiric antibiotic therapy is often initiated while awaiting the results of microbiological cultures. Oxygen therapy or mechanical ventilation may be required to maintain adequate oxygenation. Intravenous fluids may be necessary to maintain hydration and electrolyte balance. Adequate nutrition is essential for recovery from VAP. The development of VAP in this patient, despite being placed in the Trendelenburg position, highlights the delicate balance between the benefits and risks associated with different patient positioning strategies in the context of RPA management. While the Trendelenburg position can help reduce the risk of aspiration of pus from the retropharyngeal abscess, it may also increase the risk of VAP.<sup>14-16</sup>

In light of the potential risks associated with the Trendelenburg position, alternative positioning strategies have been explored. The lateral Trendelenburg position, where the patient is positioned on their side with the head of the bed slightly tilted downwards, has emerged as a potential compromise. This position aims to retain the benefits of gravity-assisted drainage while minimizing the risk of VAP associated with the supine Trendelenburg position. The lateral Trendelenburg position is a nuanced variation of the traditional Trendelenburg position, offering a potentially safer and more effective approach to managing certain clinical conditions. In this position, the patient is placed on their side with the head of the bed slightly tilted downwards. This seemingly simple adjustment offers several potential advantages over the supine Trendelenburg position, particularly in the context of mechanically ventilated patients. One of the most significant advantages of the lateral Trendelenburg position is its potential to reduce the risk of VAP. VAP is a serious complication of mechanical ventilation, often caused by the aspiration of oropharyngeal and gastric secretions into

the lower respiratory tract. These secretions can harbor bacteria, leading to infection and inflammation in the lungs. By positioning the patient on their side, the lateral Trendelenburg position helps to redirect these secretions away from the trachea and towards the dependent side of the mouth and pharynx. This reduces the likelihood of aspiration and subsequent VAP development. The lateral Trendelenburg position can also enhance lung mechanics by promoting better ventilation and perfusion matching. Ventilation refers to the movement of air in and out of the lungs, while perfusion refers to the blood flow through the lungs. In the supine Trendelenburg position, the lower parts of the lungs may be compressed, leading to decreased ventilation. Additionally, blood flow tends to be greater in the lower parts of the lungs due to gravity. This mismatch between ventilation and perfusion can result in less efficient gas exchange and potentially compromise oxygenation. The lateral Trendelenburg position helps to alleviate this mismatch by redistributing both air and blood flow within the lungs. This can lead to improved oxygenation and reduced work of breathing, particularly in patients with respiratory compromise. The lateral Trendelenburg position may also offer increased comfort for patients compared to the supine Trendelenburg position. By distributing pressure more evenly across the body, this position can reduce the risk of pressure sores, especially in patients with limited mobility or those who require prolonged bed rest. Furthermore, the lateral Trendelenburg position may feel more natural and less restrictive for some patients, promoting a sense of comfort and well-being. As discussed earlier, this position can help minimize the risk of aspiration of pus in RPA while reducing the likelihood of VAP. In patients with OHS, the lateral Trendelenburg position can improve lung function and gas exchange, potentially reducing the need for mechanical ventilation. This position may be beneficial during post-operative recovery, particularly after surgeries involving the neck or thorax, to promote drainage and enhance lung function. In addition to the lateral Trendelenburg position, other

alternative positioning strategies have been explored to optimize patient care and minimize the risks associated with prolonged bed rest and mechanical ventilation. These strategies offer various benefits and can be tailored to individual patient needs and clinical scenarios. The semi-recumbent position, in which the head of the bed is elevated to 30-45 degrees, is a widely used position for mechanically ventilated patients. It is often considered the standard of care in many intensive care units (ICUs) due to its versatility and potential benefits. The semi-recumbent position helps to reduce the risk of aspiration by utilizing gravity to keep stomach contents and oral secretions from entering the lower respiratory tract. This is particularly important in patients with impaired swallowing or those receiving enteral feeding. Elevating the head of the bed can also improve lung function by facilitating better chest expansion and diaphragm movement. This can lead to improved ventilation and oxygenation. The semi-recumbent position can be more comfortable for patients than lying completely flat, as it allows for better head and neck alignment and reduces pressure on the back. Continuous lateral rotation therapy (CLRT) involves the use of a specialized bed that continuously rotates the patient from side to side. CLRT can help to improve lung mechanics by promoting better ventilation and perfusion matching throughout the lungs. This can lead to improved oxygenation and reduced work of breathing. By continuously changing the patient's position, CLRT helps to prevent the pooling of secretions in any one area of the lungs, reducing the risk of bacterial colonization and VAP development. The constant rotation also helps to redistribute pressure across the body, minimizing the risk of pressure sores, especially in patients with limited mobility or those who require prolonged bed rest. CLRT can also aid in the drainage of secretions from the lungs and other areas, such as the abdomen or surgical sites. The choice of positioning strategy should be individualized based on the patient's specific needs and clinical condition. The patient's underlying health condition, such as respiratory

failure, neurological impairment, or cardiovascular disease, can influence the choice of positioning strategy. The severity of the patient's illness and the presence of any complications, such as sepsis or multi-organ dysfunction, can also guide the decision. The risk of complications, such as VAP, pressure sores, or hemodynamic instability, should be carefully assessed when choosing a positioning strategy. The patient's comfort and preferences should also be taken into consideration. The optimal positioning strategy for a given patient will depend on a variety of factors, including the patient's underlying health condition, the severity of their illness, and the presence of any complications. It is important to individualize the positioning strategy based on the patient's specific needs and goals.<sup>17,18</sup>

The management of retropharyngeal abscess (RPA) often presents a complex clinical challenge, requiring careful consideration of various factors to ensure optimal patient outcomes. One of the key aspects of RPA management is determining the optimal patient positioning strategy. The decision regarding the ideal position should be individualized, taking into account the unique circumstances of each patient. The patient's overall clinical condition, including their age, comorbidities, and severity of illness, should be carefully considered. Elderly patients or those with multiple comorbidities may be more vulnerable to complications associated with certain positions. The size and location of the abscess can influence the choice of positioning strategy. Larger abscesses or those located closer to the airway may necessitate positions that promote better drainage and minimize the risk of aspiration. The presence of complications, such as airway compromise, sepsis, or mediastinal extension, can significantly impact the decision-making process. Positions that optimize airway management and prevent further complications should be prioritized. If the patient requires mechanical ventilation, the risk of VAP should be carefully assessed. Certain positions, such as the Trendelenburg position, may increase the risk of VAP, while others, such as the lateral Trendelenburg

position, may offer a lower risk profile. The patient's hemodynamic stability, including their blood pressure and heart rate, should be monitored when considering different positions. Some positions may affect blood flow and cardiac function, potentially leading to hemodynamic instability. The patient's neurological status should also be taken into account. If the patient has any neurological deficits or is at risk of increased intracranial pressure, certain positions may need to be avoided or modified. While prioritizing patient safety and optimizing clinical outcomes, patient comfort should also be considered. Prolonged positioning in certain postures can lead to discomfort or pressure sores. Regular repositioning and pressure relief measures should be implemented to enhance patient comfort. A comprehensive assessment of the potential benefits and risks associated with each position is crucial for informed decision-making. Healthcare providers should carefully consider the individual patient's needs and clinical condition, weighing the potential advantages and disadvantages of each position. A multidisciplinary approach, involving otolaryngologists, intensivists, and nursing staff, is often necessary to ensure that all aspects of the patient's care are considered when making decisions about positioning. This collaborative approach can help to optimize patient outcomes and minimize complications.<sup>19,20</sup>

#### 4. Conclusion

This case report highlights the complexities of managing a retropharyngeal abscess (RPA), particularly in the context of a patient with poorly controlled diabetes who developed ventilator-associated pneumonia (VAP) despite being placed in the Trendelenburg position. While the Trendelenburg position has traditionally been used to minimize aspiration risk in RPA, this case underscores the potential for VAP development and the need for careful consideration of alternative positioning strategies, such as the lateral Trendelenburg position. The management of RPA requires a multidisciplinary approach, involving otolaryngologists, intensivists,

and other healthcare professionals. Individualized patient care, taking into account the patient's overall clinical condition, the size and location of the abscess, the presence of complications, and the risk of VAP, is crucial for informed decision-making and optimizing patient outcomes.

#### 5. References

1. Swain SK, Sahu MC. Retropharyngeal abscess leading to fatal airway obstruction in a child – A case report. *Pediatr Pol.* 2016; 91(4): 370–3.
2. Hari MS, Nirvala KD. Retropharyngeal abscess presenting with upper airway obstruction. *Anaesthesia.* 2003; 58(7): 714–5.
3. Alkan S, Koşar AT, Ozkaya I, Dadaş B. Coexistence of laryngeal mucormycosis with retropharyngeal abscess causing acute upper airway obstruction. *J Otolaryngol Head Neck Surg.* 2008; 37(3): E73-5.
4. Rao MS, Linga Raju Y, Vishwanathan P. Anaesthetic management of difficult airway due to retropharyngeal abscess. *Indian J Anaesth.* 2010; 54(3): 246–8.
5. Borgohain B. Prompt restoration of airway along with rapid neurological recovery following ultrasonography-guided needle aspiration of a tubercular retropharyngeal abscess causing airway obstruction. *Singapore Med J.* 2011; 52(11): e229-31.
6. Lynn E, Ping T, Keng Y, Singh R, Kwong W, Soon T, et al. Retropharyngeal abscess - A complication of laryngeal mask airway. *J Surg Case Rep.* 2012; 2012(10): 7.
7. Choudhuri AH, Kumar M. Airway complication due to retropharyngeal spread of epidural abscess during prone position. *J Anaesthesiol Clin Pharmacol.* 2014; 30(1): 110–1.
8. Al-Naami AQ, Ali Khan L, Ali Athlawy Y, Sun Z. *Ochrobactrum anthropi* induced retropharyngeal abscess with mediastinal extension complicating airway obstruction: a

- case report. *J Med Radiat Sci.* 2014; 61(2): 126–9.
9. Ayyash R, Pradhan C, Syed MA. Airway management in a rare case of tuberculous retropharyngeal abscess with cervical spine instability. *Anaesthesia Cases.* 2015; 3(1): 64–8.
  10. Samanta S, Samanta S, Aggarwal R, Soni KD. Airway compromise during central venous cannulation in an undiagnosed tubercular retropharyngeal abscess: a case report. *Ann Card Anaesth.* 2015; 18(4): 596–8.
  11. Hu X, Liu L. A huge retropharyngeal abscess causing airway and esophageal obstruction associated with cervical spine tuberculosis. *Spine J.* 2016; 16(4): e227-9.
  12. LeRiger M, Miler V, Tobias J, Raman VT, Elmaraghy C, Jatana K. Potential for severe airway obstruction from pediatric retropharyngeal abscess. *Int Med Case Rep J.* 2017; 10: 381–4.
  13. Lin J, Wu X-M, Feng J-X, Chen M-F. Retropharyngeal abscess presenting as acute airway obstruction in a 66-year-old woman: a case report. *World J Clin Cases.* 2019; 7(22): 3838–43.
  14. Weldetsadik AY, Bedane A, Riedel F. Retropharyngeal tuberculous abscess: A rare cause of upper airway obstruction and obstructive sleep apnea in children: a case report. *J Trop Pediatr.* 2019; 65(6): 642–5.
  15. Thota A, Sillo MO, Gangu K, Manzoor K. Peri-Tonsillar Abscess with Retropharyngeal Extension Leading to Airway Narrowing and Concomitant Pre-Septal Cellulitis. In: *D48 CRITICAL CARE CASE REPORTS: INFECTION AND SEPSIS II.* American Thoracic Society. 2019.
  16. Inman BL, Bridwell RE, Larson NP, Goss S, Oliver J. Retropharyngeal abscess with severe airway compromise following anterior cervical spine surgery: a case report. *Cureus.* 2021; 13(12): e20754.
  17. Da’Costa A, Osonuga A, Howard T, Johnston C, Osonuga A, Okoye G. Early recognition of airway compromise following a retropharyngeal abscess in the emergency department. *Vis J Emerg Med.* 2022; 26(101231): 101231.
  18. Chettri M, Das S. Acute retropharyngeal abscess in a child: a potential threat for airway compromise. *Vis J Emerg Med.* 2022; 27(101279): 101279.
  19. Hehsan MR, Muhd Shukeri WFW, Chettiar RSRMS, Abidin HZ. Successful airway management in a patient with left parapharyngeal and retropharyngeal abscess with mediastinal extension during COVID-19 pandemic. *Anaesth Pain Intensive Care.* 2022; 26(1): 128–9.
  20. Hanna JJ, Guastadisegni JM, Kouma MA, Knez EB, Arasaratnam RJ, Storey DF. Blastomycosis presenting with acute airway obstruction from a retropharyngeal abscess and complicated by severe hypokalemia during posaconazole therapy: a case report and review of literature. *Open Forum Infect Dis.* 2022; 9(8): ofac414.