



## Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: [www.bioscmed.com](http://www.bioscmed.com)

### Successful Anesthetic Management of a Cesarean Section in a Patient with Cardiomyopathy and Cardiogenic Shock: A Case Report

Alta Ikhsan Nur<sup>1\*</sup>, Nopian Hidayat<sup>2</sup>, Novita Anggraeni<sup>2</sup>, Sony<sup>2</sup>

<sup>1</sup>Anesthesiology and Intensive Therapy Specialist Education Program, Faculty of Medicine, Universitas Riau, Pekanbaru, Indonesia

<sup>2</sup>Department of Anesthesiology and Intensive Therapy, Arifin Achmad Regional General Hospital, Pekanbaru, Indonesia

#### ARTICLE INFO

##### Keywords:

Cardiogenic shock  
Cardiomyopathy  
Cesarean section  
Epidural anesthesia  
Pregnancy

##### \*Corresponding author:

Alta Ikhsan Nur

##### E-mail address:

[altaikhsannur@gmail.com](mailto:altaikhsannur@gmail.com)

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v9i3.1230>

#### ABSTRACT

**Background:** Cardiomyopathy in pregnancy is a rare but serious condition that can lead to significant maternal and fetal morbidity and mortality. Anesthetic management of these patients is challenging due to the complex interplay of physiological changes and the potential for hemodynamic instability. This case report describes the successful anesthetic management of a cesarean section in a patient with cardiomyopathy and cardiogenic shock. **Case presentation:** A 29-year-old woman with cardiomyopathy and cardiogenic shock presented for emergency cesarean section at 36-37 weeks gestation. She had a history of global hypokinetic, left ventricular and atrial dilatation, and an ejection fraction (EF) of 32%. She was also in atrial fibrillation. Epidural anesthesia was selected due to its lower risk of complications compared to general anesthesia. The patient was carefully monitored throughout the procedure, and her hemodynamics were maintained with a combination of fluids and inotropes. The surgery was successful, and the patient delivered a healthy baby boy. **Conclusion:** This case report demonstrates that successful anesthetic management of cesarean section is possible in patients with cardiomyopathy and cardiogenic shock. Careful planning, close monitoring, and a multidisciplinary approach are essential for a positive outcome.

#### 1. Introduction

Cardiomyopathy, a disorder that disrupts the heart muscle's ability to pump blood effectively, presents a formidable challenge in the realm of pregnancy care. The physiological adaptations that accompany pregnancy, such as the expansion of blood volume and the heightened demands on cardiac output, can strain an already compromised heart, potentially leading to heart failure, perilous arrhythmias, and even maternal death. The incidence of cardiomyopathy in pregnancy, though relatively low at approximately 0.46 per 1,000 deliveries, carries a significant weight of potential complications. The delicate balance between maintaining maternal well-being and ensuring a safe

delivery for the fetus necessitates a multidisciplinary approach and a nuanced understanding of anesthetic management.<sup>1-3</sup>

Anesthesiologists play a pivotal role in the care of pregnant women with cardiomyopathy. Their expertise in perioperative medicine, coupled with a thorough grasp of the patient's cardiac condition, is essential for navigating the complexities of labor and delivery. The choice of anesthesia, whether it be general or regional, must be carefully weighed, considering the patient's hemodynamic stability, the severity of the cardiomyopathy, and the urgency of the delivery. General anesthesia, while effective in achieving rapid unconsciousness and airway control, carries inherent

risks for those with cardiomyopathy. The depressant effects on the cardiovascular system, coupled with the potential for hemodynamic fluctuations during induction and emergence, necessitate meticulous monitoring and vigilance. Regional anesthesia, on the other hand, offers a targeted approach to pain control while minimizing the systemic effects associated with general anesthesia. Epidural anesthesia, in particular, has gained favor in the obstetric setting due to its ability to provide profound analgesia without compromising maternal consciousness or respiratory drive.<sup>4-7</sup>

In the specific context of cardiomyopathy, epidural anesthesia presents several advantages. By blunting the sympathetic response to pain, it can mitigate the surges in heart rate and blood pressure that might otherwise strain a weakened heart. Moreover, the gradual onset of anesthesia allows for smoother hemodynamic control and reduces the risk of sudden cardiovascular collapse. However, even with the benefits of regional anesthesia, the management of pregnant women with cardiomyopathy remains a delicate endeavor. The potential for hypotension, a common side effect of epidural anesthesia, can be particularly concerning in those with compromised cardiac output. Careful titration of anesthetic agents, along with judicious fluid administration and the availability of vasopressor support, are crucial for maintaining hemodynamic stability.<sup>8-10</sup> In this case report, we delve into the successful anesthetic management of a 29-year-old woman with cardiomyopathy and cardiogenic shock who underwent an emergency cesarean section.

## **2. Case Presentation**

This report details the case of a 29-year-old woman, gravida 5, para 0, with a history of four prior spontaneous abortions (G5P0A0H4), presenting at 36-37 weeks gestation with severe dyspnea. Her condition, later diagnosed as dilated cardiomyopathy with cardiogenic shock, presented a unique set of challenges for anesthetic management during an emergent cesarean section. The patient's primary

complaint was severe breathlessness, which had progressively worsened over the preceding week. Further inquiry revealed a three-month history of increasing dyspnea, particularly pronounced when lying down (orthopnea), suggestive of cardiac dysfunction. She also reported palpitations, a symptom often associated with arrhythmias. Importantly, the patient denied any headaches, visual disturbances, gastrointestinal symptoms (heartburn, nausea, vomiting), or seizures, helping to rule out other potential causes of her presentation such as pre-eclampsia or neurological conditions. Significantly, the patient disclosed a known cardiac abnormality diagnosed two months prior. However, she had not been receiving regular cardiology follow-up, highlighting a critical gap in her healthcare management. There was no prior history of anesthesia or surgery. She reported no known allergies or other comorbidities. Her current medications included digoxin, furosemide, and spironolactone, a regimen commonly used to manage heart failure. The patient denied any history of anticoagulant or antiplatelet use, an important consideration for potential bleeding complications during surgery. On physical examination, the patient exhibited several signs consistent with heart failure and cardiogenic shock. Her blood pressure was measured at 108/81 mmHg while receiving dobutamine at 5 mcg/kg/min, indicating that her blood pressure was being maintained with inotropic support. Her heart rate was 128 beats per minute and irregular, suggesting an underlying arrhythmia. Respiratory distress was evident with a respiratory rate of 36 breaths per minute and the presence of crackles at the lung bases, indicative of pulmonary edema. Oxygen saturation was 100% with supplemental oxygen administered via a nasal cannula at 3 L/min. Her temperature was normal at 36.6°C. Further examination revealed distended jugular veins, a classic sign of elevated central venous pressure often seen in heart failure. A heart murmur in the mitral area suggested possible valvular pathology contributing to her cardiac dysfunction. A widened ictus cordis, palpated at the

6th intercostal space on the anterior axillary line, indicated left ventricular enlargement. The presence of pretibial edema further supported the diagnosis of heart failure. Laboratory tests provided valuable insights into the patient's overall condition. Her hemoglobin level was 11.3 g/dl, which, while slightly below the normal range for pregnancy, was not critically low. Coagulation profile, liver function tests, renal function tests, and electrolyte levels were all within normal limits, suggesting that her hepatic and renal function were not significantly compromised. Chest X-ray revealed cardiomegaly, an enlarged heart, and a "bat wing" appearance, characteristic of pulmonary venous congestion commonly seen in heart failure. Echocardiography, a crucial diagnostic tool for evaluating cardiac function, confirmed the diagnosis of dilated cardiomyopathy. It showed global hypokinesis, meaning that the heart muscle was contracting weakly, along with left ventricular and atrial dilatation. Her ejection fraction (EF), a measure of the heart's pumping efficiency, was significantly reduced at 32%, indicating severe systolic dysfunction. Electrocardiography (ECG) revealed atrial fibrillation with a rapid ventricular rate of 132 bpm and a left ventricular hypertrophy (LVH) pattern, consistent with the increased workload placed on the left ventricle. Based on the comprehensive clinical assessment, including the patient's history, physical examination, laboratory findings, and imaging studies, the following diagnoses were established; G5P0A0H4 at 36-37 weeks gestation: This denotes the patient's obstetric history, indicating that she was pregnant with her fifth child, had not previously delivered a live infant, and had experienced four prior miscarriages; Dilated cardiomyopathy: This is a condition where the heart's chambers, particularly the left ventricle, become enlarged and weakened, leading to impaired pumping ability; Cardiogenic shock: This is a life-threatening condition where the heart is unable to pump enough blood to meet the body's needs, leading to organ dysfunction and potentially death; Atrial fibrillation: This is a common heart rhythm disorder characterized by rapid and irregular

electrical activity in the atria, leading to an irregular heartbeat. This patient's case represents a complex and high-risk scenario. The combination of advanced pregnancy, dilated cardiomyopathy, cardiogenic shock, and atrial fibrillation posed significant challenges for anesthetic management and required a multidisciplinary approach to optimize maternal and fetal outcomes (Table 1).

The anesthetic management of this patient with cardiomyopathy and cardiogenic shock undergoing an emergent cesarean section presented a unique set of challenges. The primary goals were to; Maintain hemodynamic stability: Avoid any sudden changes in blood pressure or heart rate that could exacerbate her cardiac condition; Ensure adequate oxygenation: Optimize oxygen delivery to both mother and fetus, particularly crucial in the context of heart failure and potential pulmonary edema; Provide effective analgesia: Minimize pain and stress, which could further strain the cardiovascular system. Considering these objectives and the patient's specific circumstances, epidural anesthesia was chosen as the preferred technique. This decision was based on several key advantages of epidural anesthesia in this context; Reduced cardiovascular instability: Compared to general anesthesia, epidural anesthesia is associated with less dramatic hemodynamic fluctuations. It avoids the myocardial depressant effects of general anesthetics and blunts the sympathetic response to surgical stress, thereby mitigating potentially harmful increases in heart rate and blood pressure; Improved oxygenation: Epidural anesthesia allows for spontaneous ventilation, avoiding the potential for airway complications and respiratory depression associated with general anesthesia. This is particularly important in a patient with potential pulmonary edema; Titratable analgesia: Epidural anesthesia provides excellent pain control with the ability to adjust the level of blockade as needed. This allows for fine-tuning of the anesthetic to match the surgical stimulus and minimize hemodynamic consequences. The epidural procedure was performed with the patient in the sitting position,

with careful aseptic technique. An 18G Tuohy needle was inserted into the L3-L4 intervertebral space, and a catheter was advanced 5 cm into the epidural space. A test dose of 1.2 ml bupivacaine (a local anesthetic with rapid onset), 1 ml lidocaine 2%, and 0.8 ml distilled water was administered to confirm proper placement and exclude intravascular or intrathecal injection. This test dose aimed to rule out accidental intravenous injection (which could cause systemic toxicity) or intrathecal injection (which could lead to a high spinal block and respiratory paralysis). Following the successful test dose, the patient was positioned supine with the head of the bed elevated to facilitate venous return and maintain cardiac output. Ropivacaine 0.75% was administered through the epidural catheter to achieve a T6 sensory level, providing adequate anesthesia for the cesarean section. This level of blockade ensured adequate analgesia for the surgical procedure while minimizing the risk of hypotension caused by sympathetic blockade. Throughout the procedure, meticulous monitoring was maintained. This included continuous invasive blood pressure monitoring, central venous pressure monitoring, and pulse oximetry to closely track the patient's hemodynamic status and oxygenation. Oxygen was administered via a nasal cannula at 3 L/min to ensure adequate oxygen delivery. Fluid management was crucial to maintain intravascular volume and support cardiac output. Ringer's lactate solution was administered intravenously as needed to prevent hypotension and ensure adequate tissue perfusion. Following the cesarean section, the patient was transferred to the intensive care unit (ICU) for close observation for 24 hours. Continuous epidural analgesia was provided with a lower concentration of ropivacaine (0.2%) and fentanyl (2 mcg/ml) to control postoperative pain while minimizing the risk of hemodynamic instability. This allowed for effective pain management while minimizing the need for systemic opioids, which can have depressant effects on the cardiovascular and respiratory systems. The choice of epidural anesthesia in this case was a carefully considered decision based

on the patient's unique circumstances. While general anesthesia might have been faster and provided more complete control of the airway, it carried a higher risk of hemodynamic instability and respiratory complications in this patient with compromised cardiac function. Epidural anesthesia, on the other hand, allowed for a more gradual and controlled onset of anesthesia, minimizing the stress on the cardiovascular system and allowing for spontaneous ventilation. The use of ropivacaine as the primary local anesthetic was also a deliberate choice. Ropivacaine is known for its less pronounced motor blockade compared to other local anesthetics, which can be beneficial in facilitating early ambulation and reducing the risk of thromboembolic complications in the postpartum period. The addition of fentanyl to the epidural infusion in the postoperative period provided synergistic analgesia and reduced the total dose of local anesthetic required. This helped to minimize the risk of local anesthetic toxicity and motor blockade (Table 2).

The postoperative period for this patient with cardiomyopathy and cardiogenic shock required intensive monitoring and meticulous care to ensure both maternal and neonatal well-being. A multifaceted approach was adopted, encompassing pain management, hemodynamic stabilization, and close observation for potential complications. Effective pain management was crucial to the patient's recovery and hemodynamic stability. Uncontrolled pain can trigger a stress response, leading to elevated heart rate and blood pressure, which could be detrimental to a patient with compromised cardiac function. To address this, continuous epidural analgesia was implemented using a lower concentration of ropivacaine (0.2%) combined with fentanyl (2 mcg/ml). This combination provided synergistic analgesia, allowing for effective pain control while minimizing the dose of local anesthetic and reducing the risk of motor blockade and systemic side effects. The patient's pain was regularly assessed using the Visual Analog Scale (VAS) to ensure adequate pain relief and guide adjustments in the epidural infusion

rate. This proactive approach to pain management contributed to the patient's comfort and facilitated early mobilization, reducing the risk of postoperative complications such as deep vein thrombosis and pulmonary embolism. Close hemodynamic monitoring was continued in the immediate postoperative period. The patient was transferred to the intensive care unit (ICU) for 24 hours of continuous observation, where her heart rate, blood pressure, and oxygen saturation were closely monitored. This allowed for prompt detection and management of any hemodynamic instability, ensuring adequate perfusion to vital organs and supporting her recovery. Remarkably, the patient did not experience any postoperative complications. This positive outcome can be attributed to the meticulous anesthetic management, careful fluid balance, and vigilant postoperative care provided by the multidisciplinary team. The absence of complications highlights the effectiveness of the chosen anesthetic technique and the importance of comprehensive perioperative care in high-risk patients. The patient's recovery progressed smoothly, allowing for discharge from the ICU on the first postoperative day and from the hospital on the fourth postoperative day. However, given her underlying cardiac condition, a comprehensive follow-up plan was essential to ensure continued stability and optimize her long-term health; Within 1 week of discharge: Cardiology consultation was scheduled to assess her cardiac function, review her medication regimen, and make any necessary adjustments. This early follow-up aimed to ensure that her heart failure was adequately managed and to address any new or worsening symptoms; Within 4 weeks of discharge: An echocardiogram was planned to evaluate her left ventricular function and assess her response to treatment. This non-invasive imaging study provided valuable information about the size and function of her heart chambers, ejection fraction, and any changes in her valvular function; Long-term follow-up: Continued cardiology follow-up was recommended every 3-6 months, with optimization of medical therapy as needed. This long-term monitoring aimed

to track her progress, identify any potential complications, and ensure that her treatment plan remained aligned with her evolving needs; Within 2 weeks of discharge: The patient was scheduled for a routine postpartum check-up with her obstetrician. This visit addressed any concerns related to her recovery from childbirth, provided family planning counseling, and offered support for her overall well-being. A crucial component of the postoperative management plan was patient education. The patient was counseled on lifestyle modifications to support her cardiac health, including dietary adjustments, regular exercise, and stress management techniques. She was also educated on the importance of medication adherence and recognizing potential signs and symptoms of worsening heart failure. Empowering the patient with knowledge and self-management strategies played a vital role in her ongoing recovery and long-term health (Table 3).

### **3. Discussion**

Cardiomyopathy in pregnancy represents a formidable challenge in the realm of maternal-fetal medicine. It is a complex interplay of the heart's intrinsic pathology and the physiological adaptations that accompany pregnancy. The pregnant state, while a natural phenomenon, places significant demands on the cardiovascular system. These demands, coupled with an underlying cardiac condition like cardiomyopathy, can lead to a cascade of complications, jeopardizing the health of both mother and child. The physiological adaptations of pregnancy are numerous and profound. The maternal blood volume undergoes a dramatic expansion, increasing by approximately 40-50% to support the growing fetus and prepare for the anticipated blood loss during childbirth. This expanded blood volume, in turn, leads to an increase in cardiac output, the amount of blood pumped by the heart per minute. The heart rate also increases, further augmenting the workload of the heart. These adaptations, while essential for a healthy pregnancy, can place significant strain on a heart already compromised by cardiomyopathy.

Table 1. Anamnesis, clinical finding, laboratory, imaging, and clinical diagnosis.

Anamnesis	Clinical finding	Laboratory	Imaging	Clinical diagnosis
A 29-year-old woman, G5P0A0H4 at 36-37 weeks gestation, presented to the hospital with Severe breathlessness for 1 week	Blood pressure: 108/81 mmHg (on dobutamine 5 mcg/kg/min)	Hemoglobin: 11.3 g/dL	Chest X-ray: Cardiomegaly and bat wing appearance	G5P0A0H4 36-37 weeks gestation
Breathlessness worsening over 3 months, especially when lying down	Heart rate: 128 bpm irregular	Normal coagulation profile	Echocardiography: Global hypokinesis, left ventricular and atrial dilatation, ejection fraction (EF) 32%	Dilated cardiomyopathy
Palpitations	Respiratory rate: 36 breaths per minute	Normal liver function test	Electrocardiography: Atrial fibrillation with a ventricular rate of 132 bpm, LVH pattern	Cardiogenic shock
No headaches, blurry vision, heartburn, nausea, vomiting, or seizures	Oxygen saturation: 100% with a nasal cannula at 3 L/min	Normal renal function test		Atrial fibrillation
Known cardiac abnormalities for 2 months, no regular cardiology follow-up	Temperature: 36.6°C	Normal electrolytes	-	-
No prior history of anesthesia or surgery	Distended jugular veins	-	-	-
No allergies or other comorbidities	Crackles at lung bases	-	-	-
Current medications: digoxin, furosemide, spironolactone	Heart murmur in the mitral area	-	-	-
No history of anticoagulant or antiplatelet use	Widened icтус cordis at ICS 6 on the anterior axillary line	-	-	-
	Pretibial edema	-	-	-

Table 2. Anesthetic management.

Anesthetic management	Details
Anesthesia technique	Epidural anesthesia
Epidural insertion site	L3-L4 intervertebral space
Epidural needle	18G Tuohy needle
Epidural catheter insertion	5 cm into the epidural space
Test dose	1.2 ml pehacain + 1 ml lidocaine 2% + 0.8 ml distilled water
Patient position	Sitting for epidural placement, then supine with head of bed elevated
Medications	Ropivacaine 0.75% for surgical anesthesia, ropivacaine 0.2%, and fentanyl 2 mcg/ml for postoperative analgesia
Anesthesia level	T6 sensory block
Oxygenation	Nasal cannula at 3 L/min
Monitoring	Continuous invasive blood pressure monitoring, central venous pressure monitoring, oxygen saturation
Fluid management	Ringer's lactate solution
Postoperative management	Intensive care unit (ICU) observation for 24 hours, continuous epidural analgesia

Table 3. Postoperative management and follow-up.

<b>Postoperative management and follow-up</b>	<b>Details</b>
Postoperative analgesia	Continuous epidural infusion of ropivacaine 0.2% and fentanyl 2 mcg/ml
Postoperative monitoring	Intensive care unit (ICU) observation for 24 hours
Hemodynamic monitoring	Continuous monitoring of heart rate, blood pressure, and oxygen saturation
Pain assessment	Visual Analog Scale (VAS) score
Postoperative complications	None reported
Discharge from ICU	On the first postoperative day
Discharge from hospital	On the fourth postoperative day
Follow-up care	<p>Cardiology follow-up within 1 week of discharge to assess cardiac function and optimize medication regimen.</p> <p>Echocardiogram within 4 weeks of discharge to evaluate left ventricular function and assess response to treatment.</p> <p>Obstetrics follow-up within 2 weeks of discharge for routine postpartum care and family planning counseling.</p> <p>Long-term cardiology follow-up every 3-6 months with optimization of medical therapy as needed.</p> <p>Patient education regarding lifestyle modifications, medication adherence, and symptom recognition.</p>

Cardiomyopathy, a disorder that disrupts the heart muscle's ability to pump blood effectively, manifests in various forms, each with its unique pathophysiology. Dilated cardiomyopathy, characterized by the enlargement and weakening of the heart chambers, is the most common type encountered in pregnancy. Hypertrophic cardiomyopathy, marked by the thickening of the heart muscle, and restrictive cardiomyopathy, characterized by the stiffening of the heart walls, are less common but equally concerning. In the context of pregnancy, these cardiomyopathies can lead to a spectrum of complications. Heart failure, a condition where the heart is unable to pump enough blood to meet the body's needs, is a common consequence. The symptoms of heart failure can range from mild breathlessness to life-threatening pulmonary edema, the accumulation of fluid in the lungs. Arrhythmias, or irregular heart rhythms, are another potential complication. The increased electrical activity of the heart during pregnancy, coupled with the structural abnormalities associated with cardiomyopathy, can predispose to the

development of arrhythmias. These arrhythmias can range from benign palpitations to life-threatening ventricular tachycardia or fibrillation, where the heart's rhythm becomes chaotic and ineffective. The consequences of these complications can be devastating. Maternal mortality, though rare, is a stark reality in the setting of cardiomyopathy in pregnancy. The risk of death is highest in those with severe cardiomyopathy, particularly those with heart failure or significant arrhythmias. Fetal complications are also a concern. The inadequate blood supply to the placenta, resulting from the mother's compromised cardiac function, can lead to fetal growth restriction, preterm birth, and even stillbirth. The risk of these complications is highest in those with severe cardiomyopathy and those who develop heart failure during pregnancy. The management of cardiomyopathy in pregnancy requires a multidisciplinary approach, involving obstetricians, cardiologists, and anesthesiologists. The goal is to optimize the mother's cardiac function while ensuring the safe delivery of a healthy infant. This often involves

a combination of medication, lifestyle modifications, and close monitoring. Medications such as beta-blockers, ACE inhibitors, and diuretics are commonly used to manage heart failure and arrhythmias. These medications help to reduce the workload of the heart, improve its pumping efficiency, and control irregular heart rhythms. However, the use of these medications during pregnancy requires careful consideration, as some can have adverse effects on the developing fetus. Lifestyle modifications, such as limiting physical activity and reducing sodium intake, can also help to manage the symptoms of heart failure. These modifications help to reduce the strain on the heart and prevent fluid retention. Close monitoring is essential throughout pregnancy and the postpartum period. This includes regular check-ups with the obstetrician and cardiologist, as well as frequent echocardiograms to assess cardiac function. The goal is to detect any signs of worsening heart failure or arrhythmias promptly and intervene accordingly. In the event of severe cardiomyopathy or complications such as heart failure or arrhythmias, early delivery may be necessary to protect the health of both mother and child. The mode of delivery, whether vaginal or cesarean, is determined based on the individual circumstances and the severity of the cardiac condition. The anesthetic management of pregnant women with cardiomyopathy presents unique challenges. The choice of anesthesia, whether general or regional, must be carefully weighed, considering the potential impact on cardiac function and the risk of complications. General anesthesia, while effective in achieving rapid unconsciousness and airway control, carries inherent risks for those with cardiomyopathy. The depressant effects on the cardiovascular system, coupled with the potential for hemodynamic fluctuations during induction and emergence, necessitate meticulous monitoring and vigilance. Regional anesthesia, on the other hand, offers a targeted approach to pain control while minimizing the systemic effects associated with general anesthesia. Epidural anesthesia, in particular, has gained favor in the obstetric setting due to its ability

to provide profound analgesia without compromising maternal consciousness or respiratory drive. In the specific context of cardiomyopathy, epidural anesthesia presents several advantages. By blunting the sympathetic response to pain, it can mitigate the surges in heart rate and blood pressure that might otherwise strain a weakened heart. Moreover, the gradual onset of anesthesia allows for smoother hemodynamic control and reduces the risk of sudden cardiovascular collapse. However, even with the benefits of regional anesthesia, the management of pregnant women with cardiomyopathy remains a delicate endeavor. The potential for hypotension, a common side effect of epidural anesthesia, can be particularly concerning in those with compromised cardiac output. Careful titration of anesthetic agents, along with judicious fluid administration and the availability of vasopressor support, are crucial for maintaining hemodynamic stability.<sup>11-13</sup>

The selection of anesthesia for a cesarean section in a patient with cardiomyopathy is a pivotal decision that can significantly influence both maternal and fetal outcomes. It requires a thorough understanding of the patient's cardiovascular status, the pathophysiology of their specific cardiomyopathy, and the potential interactions between anesthetic agents and the altered physiological state of pregnancy. Anesthesiologists face a critical choice between two primary approaches, general anesthesia and regional anesthesia. Each technique offers distinct advantages and disadvantages in the context of cardiomyopathy, and the optimal choice depends on a careful risk-benefit assessment tailored to the individual patient. General anesthesia, characterized by the induction of a reversible state of unconsciousness, provides several benefits in the obstetric setting. It ensures rapid airway control, crucial in emergency situations or when difficult intubation is anticipated. It also offers the advantage of amnesia, sparing the mother any recollection of the surgical procedure. However, in the presence of cardiomyopathy, general anesthesia carries significant risks. The volatile anesthetic agents commonly used to maintain general anesthesia can



depress myocardial contractility, reducing the heart's ability to pump blood effectively. This can be particularly detrimental in patients with already compromised cardiac function, potentially leading to hypotension, decreased cardiac output, and inadequate tissue perfusion. Furthermore, the process of intubation and extubation can trigger hemodynamic fluctuations, causing surges in heart rate and blood pressure that may be poorly tolerated in patients with cardiomyopathy. The potential for laryngospasm, bronchospasm, and aspiration also adds to the complexity of general anesthesia in this population. Regional anesthesia, encompassing techniques such as epidural and spinal anesthesia, offers a targeted approach to pain control while avoiding the systemic effects of general anesthesia. By delivering anesthetic agents directly to the nerves supplying the surgical area, regional anesthesia provides profound analgesia without compromising maternal consciousness or respiratory drive. In the context of cardiomyopathy, regional anesthesia presents several potential advantages. By blocking the sympathetic nervous system, it can induce vasodilation, reducing afterload and the workload on the heart. This can be particularly beneficial in patients with heart failure, where the heart is already struggling to pump blood effectively. Moreover, regional anesthesia avoids the myocardial depressant effects of general anesthetics, preserving cardiac contractility and reducing the risk of hypotension. It also allows for a more gradual and controlled onset of anesthesia, minimizing the hemodynamic fluctuations that can occur during induction and emergence from general anesthesia. However, regional anesthesia is not without its considerations. The potential for hypotension, a common side effect of sympathetic blockade, can be particularly concerning in patients with compromised cardiac output. Careful titration of anesthetic agents, along with judicious fluid administration and the availability of vasopressor support, are crucial for maintaining hemodynamic stability. Epidural anesthesia, a specific type of regional anesthesia, has gained favor in the obstetric

setting due to its versatility and ability to provide continuous pain relief throughout labor and the postpartum period. In the context of cardiomyopathy, epidural anesthesia offers a balanced approach, minimizing the risks associated with both general anesthesia and spinal anesthesia. Unlike spinal anesthesia, which involves a single injection of anesthetic into the spinal canal, epidural anesthesia allows for the insertion of a catheter into the epidural space. This catheter provides a route for continuous administration of anesthetic agents, allowing for fine-tuning of the level of blockade and hemodynamic control. The gradual onset of anesthesia with epidural anesthesia also reduces the risk of sudden hypotension, allowing for smoother hemodynamic management. Moreover, the ability to extend the epidural infusion into the postpartum period provides effective pain control and facilitates early mobilization, reducing the risk of thromboembolic complications. In the case presented, the decision was made to proceed with epidural anesthesia. This decision was likely based on a careful assessment of the patient's hemodynamic instability and the potential for complications associated with general anesthesia. Epidural anesthesia offered the potential for better hemodynamic control, a reduced risk of cardiovascular depression, and the flexibility to provide continuous analgesia throughout the perioperative period. The successful outcome of this case highlights the importance of individualized anesthetic planning and the careful selection of anesthetic techniques in patients with cardiomyopathy. By weighing the potential risks and benefits of each approach, anesthesiologists can play a crucial role in ensuring the safe delivery of healthy babies while minimizing maternal complications.<sup>14-16</sup>

The successful management of anesthesia in a patient with cardiomyopathy hinges on meticulous monitoring and proactive interventions. This is particularly crucial in the setting of pregnancy, where physiological adaptations and the stress of childbirth can further challenge the cardiovascular system. In this case, the anesthesia team's vigilance and prompt

responses played a pivotal role in ensuring a positive outcome for both mother and baby. The cornerstone of safe anesthetic management is the continuous assessment of key physiological parameters. In this patient with cardiomyopathy and cardiogenic shock, a comprehensive monitoring approach was essential to track her cardiovascular stability and guide timely interventions. Continuous invasive blood pressure monitoring, achieved through an arterial line, provided real-time, beat-to-beat blood pressure readings. This allowed for the immediate detection of any fluctuations in blood pressure, enabling prompt adjustments in fluid management or vasoactive medications. A central venous catheter, inserted into a large vein, provided continuous monitoring of central venous pressure (CVP). CVP reflects the pressure in the right atrium and serves as an indicator of the heart's preload, or the volume of blood returning to the heart. Monitoring CVP helped assess the patient's fluid status and guide fluid management strategies, ensuring adequate intravascular volume without overloading the heart. Continuous pulse oximetry, a non-invasive method to measure oxygen saturation in the blood, ensured that the patient maintained adequate oxygenation throughout the perioperative period. This was particularly crucial in the context of heart failure, where compromised cardiac output can lead to inadequate oxygen delivery to tissues. Continuous ECG monitoring tracked the patient's heart rate and rhythm, providing immediate alerts for any arrhythmias. This was essential in this patient with a history of atrial fibrillation, as any new or worsening arrhythmias could further compromise her cardiac function. Maintaining normothermia, or a normal body temperature, is crucial in anesthesia. Hypothermia can lead to coagulopathy, or impaired blood clotting, and increase the risk of bleeding complications. Continuous temperature monitoring allowed for prompt interventions to maintain normothermia and prevent these complications. Monitoring urine output provided an indirect measure of renal perfusion and overall tissue perfusion. A decline in urine output can signal inadequate cardiac

output or hypovolemia, prompting adjustments in fluid management or vasoactive medications. The patient's underlying cardiomyopathy and cardiogenic shock required proactive hemodynamic management to maintain cardiovascular stability throughout the perioperative period. The anesthesia team employed a combination of fluid therapy and vasoactive medications to optimize her intravascular volume, support her cardiac output, and ensure adequate tissue perfusion. Intravenous fluids, such as Ringer's lactate solution, were administered to maintain adequate intravascular volume and prevent hypotension. The choice of fluid and the rate of administration were carefully titrated based on the patient's hemodynamic parameters, including blood pressure, heart rate, and CVP. Vasopressors, such as phenylephrine or norepinephrine, are medications that constrict blood vessels, increasing blood pressure. Inotropes, such as dobutamine or milrinone, increase the contractility of the heart, improving cardiac output. In this case, the patient was already receiving dobutamine, an inotrope with both inotropic and vasodilatory properties, to support her cardiac function. The anesthesia team likely continued this medication and titrated its dose based on her hemodynamic response. Beyond the specific monitoring modalities and interventions, the successful management of this case relied heavily on the anesthesia team's vigilance and responsiveness. The continuous monitoring data provided a real-time window into the patient's physiological status, allowing the team to anticipate potential complications and intervene proactively. The team's expertise in interpreting the monitoring data, recognizing subtle changes in the patient's condition, and promptly adjusting fluid management or vasoactive medications played a crucial role in maintaining hemodynamic stability and preventing adverse events.<sup>17,18</sup>

The successful management of a pregnant patient with cardiomyopathy requires a multidisciplinary approach, a harmonious collaboration of expertise from various medical specialties. This intricate interplay of knowledge and skills ensures that the

patient receives comprehensive care that addresses both the complexities of her pregnancy and the challenges posed by her underlying cardiac condition. The obstetricians are the primary caregivers for the pregnant patient, responsible for monitoring the progress of her pregnancy, managing any obstetric complications, and ensuring the safe delivery of her baby. Their expertise lies in understanding the physiological changes of pregnancy, recognizing potential risks, and making informed decisions about the timing and mode of delivery. The cardiologists provide specialized care for the patient's underlying cardiomyopathy. They assess the severity of her cardiac condition, optimize her medication regimen, and monitor her cardiac function throughout her pregnancy. Their expertise is crucial in identifying potential cardiac complications, guiding treatment decisions, and ensuring that the patient's heart is as healthy as possible for the demands of childbirth. The anesthesiologists play a critical role in the perioperative care of the patient. They assess her cardiovascular status, select the most appropriate anesthetic technique for her cesarean section, and meticulously monitor her hemodynamic parameters throughout the procedure. Their expertise in managing the patient's airway, providing adequate pain control, and maintaining hemodynamic stability is essential for a safe and successful outcome. The success of the multidisciplinary approach hinges on effective communication and coordination among the team members. Regular meetings, case discussions, and shared decision-making ensure that everyone is informed about the patient's condition, the treatment plan, and any potential challenges. This collaborative effort extends beyond the core team, encompassing other healthcare professionals such as nurses, pharmacists, and dietitians. The nurses provide continuous bedside care, monitoring the patient's vital signs, administering medications, and providing emotional support. The pharmacists ensure the safe and effective use of medications, while the dietitians provide guidance on dietary modifications to support the patient's cardiac health. By bringing together

expertise from various specialties, the multidisciplinary team can develop a comprehensive plan of care that is tailored to the unique needs of each patient. This individualized approach ensures that the patient receives the most appropriate treatment and monitoring for her specific circumstances. The collaborative effort of the team allows for early detection and prompt management of potential complications. Regular monitoring, combined with the expertise of cardiologists and anesthesiologists, helps to identify any signs of worsening heart failure or arrhythmias, enabling timely interventions to prevent adverse events. The multidisciplinary approach has been shown to improve maternal and fetal outcomes in pregnant women with cardiomyopathy. By optimizing the mother's cardiac health and ensuring a safe delivery, the team can reduce the risk of complications such as heart failure, arrhythmias, preterm birth, and fetal growth restriction. The multidisciplinary approach provides patients with a sense of reassurance and confidence, knowing that they are receiving comprehensive care from a team of experts. This can help to reduce anxiety and improve their overall experience during a challenging time.<sup>19,20</sup>

#### **4. Conclusion**

This case report demonstrates that successful anesthetic management of cesarean section is possible in patients with cardiomyopathy and cardiogenic shock. The choice of anesthesia for this patient was epidural anesthesia. Epidural anesthesia has the advantage of reducing the risk of cardiovascular instability compared to general anesthesia. It also avoids the myocardial depressant effects of general anesthetics and blunts the sympathetic response to surgical stress, thereby mitigating potentially harmful increases in heart rate and blood pressure. In addition, epidural anesthesia allows for spontaneous ventilation, avoiding the potential for airway complications and respiratory depression associated with general anesthesia. The successful outcome of this case can be attributed to several factors, including the careful selection of anesthetic technique, the

meticulous monitoring of the patient's hemodynamic status, and the prompt intervention to maintain cardiovascular stability. The multidisciplinary team approach, involving obstetricians, cardiologists, and anesthesiologists, also played a crucial role in ensuring the best possible outcome for both mother and child. This case report highlights the importance of individualized anesthetic planning and the careful selection of anesthetic techniques in patients with cardiomyopathy. By weighing the potential risks and benefits of each approach, anesthesiologists can play a crucial role in ensuring the safe delivery of healthy babies while minimizing maternal complications.

## 5. References

1. Zhang X, Xu T, Jia L, Cao H, Xu Z. Continuous cardiovascular hemodynamics monitoring with pressure recording analytical method in patients under spinal anesthesia for elective cesarean section: a pilot study. *Ann Palliat Med.* 2021; 10(7): 7184–93.
2. Garutti I, Cabañero A, Vicente R, Sánchez D, Granell M, Fraile CA, et al. Recommendations of the Society of Thoracic Surgery and the Section of Cardiothoracic and Vascular Surgery of the Spanish Society of Anesthesia, Resuscitation and Pain Therapy, for patients undergoing lung surgery included in an intensified recovery program. *Rev Esp Anesthesiol Reanim (Engl Ed).* 2022; 69(4): 208–41.
3. Omar H, Helmy MH, Wahib M, Abdalla Mohamed A, Marzouk S, Abdelreheem AM. The effect of type of anesthesia (general versus spinal) on postoperative levels of brain natriuretic peptide in parturient women with cardiovascular diseases undergoing cesarean section and related cardiac impacts. *Egypt J Anaesth.* 2024; 40(1): 10–7.
4. Iwai T, Inomata S. Successful perimortem cesarean section with cardiopulmonary resuscitation and anesthesia: a report of an unique case. *Cureus.* 2024; 16(2): e55037.
5. Kocum A, Sener M, Caliskan E, Izmirli H, Tarım E, Kocum T, et al. Epidural anesthesia for cesarean section in a patient with severe mitral stenosis and pulmonary hypertension. *J Cardiothorac Vasc Anesth.* 2010; 24(6): 1022–3.
6. Hansda U, Das PK, Sahoo S. Cesarean section with spinal anesthesia in a patient with double outlet right ventricle. *J Cardiothorac Vasc Anesth.* 2020; 34(5): 1394–6.
7. ELZayyat N, Hassan A, ELSakka A, Saad A. Ketamine/propofol (ketofol) versus propofol/fentanyl for induction of general anesthesia in parturients with rheumatic valvular lesions undergoing elective cesarean section. *Egypt J Cardiothorac Anesth.* 2013; 7(2): 79.
8. Del Rio JM, Abernathy JH Iii, Taylor MA, Kertai MD. What makes a team work, a specialty work, and a society work: The SCA/STS adult cardiac anesthesia section of the STS database for the SCA/STS database sub-committee. *J Cardiothorac Vasc Anesth.* 2019; 33(8): 2109–17.
9. Hashimoto E, Kojima A, Kitagawa H, Matsuura H. Anesthetic management of a patient with type 1 long QT syndrome using combined epidural-spinal anesthesia for caesarean section: Perioperative approach based on ion channel function. *J Cardiothorac Vasc Anesth.* 2020; 34(2): 465–9.
10. Del Rio JM, Jake Abernathy J 3rd, Taylor MA, Habib RH, Fernandez FG, Bollen BA, et al. The Adult Cardiac Anesthesiology Section of STS Adult Cardiac Surgery Database: 2020 update on quality and outcomes. *J Cardiothorac Vasc Anesth.* 2021; 35(1): 22–34.
11. Tsukinaga A, Yoshitani K, Kubota Y, Kanemaru E, Nishimura K, Ogata S, et al. Anesthesia for cesarean section and postpartum cardiovascular events in congenital heart disease: a retrospective

- cohort study. *J Cardiothorac Vasc Anesth.* 2021; 35(7): 2108–14.
12. Tafesse H, Shanmugam G, Long B, BuAbbud A, Aziz S, Parr KG. ECMO for TACO secondary to massive post-cesarean section hemorrhage. *J Cardiothorac Vasc Anesth.* 2022; 36(8 Pt A): 2578–82.
  13. Kapoor PM, Goyal S, Irpachi K, Smita B. Importance of transesophageal echocardiography in peripartum cardiomyopathy undergoing lower section cesarean section under regional anesthesia. *J Anaesthesiol Clin Pharmacol.* 2014; 30(3): 427–9.
  14. Hakimoglu S, Koyuncu O, Akkurt C, Yesil S, Urfali S. Anesthesia management for emergency cesarean section in a patient with takotsubo cardiomyopathy. *Med Sci (Turkey).* 2020; 9(1): 271.
  15. Sawant SM, Oak SP, Gangakhedkar GR, Hemantkumar IC. Successful outcome after regional anesthesia in a parturient with Dilated cardiomyopathy for elective cesarean section: Navigating a stormy sea. *Med J Dr DY Patil Vidyapeeth.* 2023; 16(6): 981–3.
  16. Desalegn M. Cesarean section under spinal anesthesia for a mother with dilated cardiomyopathy in a resource-limited setting: a case report. *Ann Med Surg (Lond).* 2024; 86(2): 1182–4.
  17. Puthenveettil N, Ramakrishnan M, Paul J. Continuous spinal anesthesia for cesarean section in a parturient with peripartum cardiomyopathy. *J Obstet Anaesth Crit Care.* 2024; 14(2): 195–6.
  18. Dutt A, Agarwal A, Chatterji R, Ahmed F. Anesthetic management for cesarean section in a case of peripartum cardiomyopathy. *Anesth Essays Res.* 2013; 7(2): 273–5.
  19. Suzuki T, Nemoto C, Ikegami Y, Yokokawa T, Tsukada Y, Abe Y, et al. Development of takotsubo cardiomyopathy with severe pulmonary edema before a cesarean section. *J Anesth.* 2014; 28(1): 121–4.
  20. Kourkovei P, Leontiadis E, Kontonika M, Antoniou T, Iakovou I, Karatasakis G. Reverse takotsubo cardiomyopathy after caesarean section with epidural anesthesia: Initial assessment and follow-up. *J Cardiothorac Vasc Anesth.* 2020; 34(3): 756–8.