



Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: www.bioscmed.com

Optimizing Sunburn Prevention in Children: A Meta-Analysis of the Efficacy of Sunscreen Application, Protective Apparel, and Shade-Seeking Behaviors

Endra Yustin Ellistasari^{1*}, Suci Widhiati¹, Sesia Pradestine¹

¹Department of Dermatology and Venereology, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia

ARTICLE INFO

Keywords:

Children
Protective clothing
Skin cancer prevention
Sunburn
Sunscreen

*Corresponding author:

Endra Yustin Ellistasari

E-mail address:

endra_yustin@yahoo.com

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v9i8.1355>

ABSTRACT

Background: Childhood sunburn significantly elevates lifelong skin cancer risk, underscoring the need for effective prevention. While sunscreen, protective apparel, and shade-seeking are advocated, a quantitative synthesis of their efficacy in children is crucial. This meta-analysis aimed to consolidate and quantify the evidence on the effectiveness of these core sun protection strategies in preventing pediatric sunburn. **Methods:** A systematic search of PubMed/MEDLINE, Scopus, Cochrane CENTRAL, and Web of Science (January 2014 - December 2024) identified randomized controlled trials and cohort studies evaluating sunscreen, protective apparel, or shade-seeking behaviors for sunburn prevention in individuals aged 0-18 years. Data on sunburn incidence were extracted, study quality assessed, and pooled Risk Ratios (RR) with 95% confidence intervals (CI) calculated using a random-effects model. **Results:** Six studies (two RCTs, four cohort studies) involving 8,500 children were included. Regular sunscreen use (SPF ≥ 30) significantly reduced sunburn incidence (RR 0.65, 95% CI 0.55-0.77). Protective apparel use also demonstrated substantial protection (RR 0.70, 95% CI 0.60-0.82). Enhanced shade provision and shade-seeking behaviors effectively lowered sunburn risk (RR 0.75, 95% CI 0.62-0.90). Multi-component strategies combining these approaches showed consistent protective benefits. **Conclusion:** This meta-analysis provides robust quantitative evidence that diligent sunscreen application, consistent use of protective apparel, and active shade-seeking are all significantly effective in reducing sunburn incidence in children. These findings strongly support multifaceted public health initiatives emphasizing comprehensive sun protection to safeguard pediatric skin health.

1. Introduction

Childhood and adolescence represent crucial developmental phases marked by significant cumulative exposure to solar ultraviolet radiation (UVR), with a substantial portion of an individual's lifetime UVR dose often acquired during these formative years. Sunburn, an acute cutaneous inflammatory response to excessive UVR, is a prevalent yet largely preventable condition in pediatric populations. While the immediate effects of sunburn include pain, erythema, and blistering, of greater concern are the long-term sequelae. A robust body of epidemiological research has consistently established a strong association between sunburn episodes in

early life and an increased risk of developing skin cancers, including cutaneous melanoma and non-melanoma skin cancers, in adulthood. Solar UVR is unequivocally classified as a Group 1 carcinogen by the International Agency for Research on Cancer. Consequently, mitigating excessive UVR exposure, particularly during the vulnerable pediatric years, is a critical public health objective. Children's skin possesses unique anatomical and physiological characteristics that render it more susceptible to UVR-induced damage compared to adult skin. These include a thinner epidermis and stratum corneum, which may allow for greater UVR penetration, and a developing melanin system that provides less intrinsic

photoprotection, especially in individuals with fairer skin types. Furthermore, DNA repair mechanisms in children, while functional, might be more easily overwhelmed by extensive UV damage, increasing the potential for mutations that can initiate carcinogenesis. Compounding this physiological vulnerability are behavioral patterns common in childhood, such as extended periods of outdoor activity, often during hours of peak UVR intensity (typically 10:00 a.m. to 4:00 p.m.), without consistent or adequate protection. Studies indicate that a significant percentage of children experience at least one sunburn annually, highlighting an ongoing need for effective and consistently applied sun protection strategies.¹⁻⁴

In response to this public health challenge, leading health organizations worldwide advocate for a multifaceted approach to sun protection for children. This typically involves a triad of primary behaviors: the diligent application of broad-spectrum sunscreen with a sun protection factor (SPF) of 30 or higher; wearing photoprotective apparel, including hats and UV-blocking sunglasses; and actively seeking or utilizing shade, especially during midday hours. Sunscreen application is a cornerstone of photoprotection, containing UV filters that absorb, reflect, or scatter UVR, thereby reducing its penetration into the skin. Broad-spectrum formulations protect against both UVB, the primary cause of sunburn and direct DNA damage, and UVA, which contributes to photoaging, indirect DNA damage, and carcinogenesis. However, the real-world effectiveness of sunscreen is often compromised by suboptimal application practices, such as using insufficient amounts or infrequent reapplication. Protective apparel offers a physical barrier against UVR and is considered a highly reliable method of sun protection. The ultraviolet protection factor (UPF) of a fabric quantifies its ability to block UVR, with factors like weave, fiber type, and color influencing its protective capacity. Wide-brimmed hats and UV-protective sunglasses are essential components, shielding highly exposed and vulnerable areas such as the face, ears, neck, and eyes. Shade-

seeking, whether utilizing natural or artificial shade, can significantly reduce direct UVR exposure, especially during peak hours. Environmental modifications to increase shade in schools and playgrounds represent important public health interventions. While shade mitigates direct UVR, some exposure to diffuse or reflected UVR can still occur, emphasizing the need for shade to be part of a comprehensive strategy. While the individual benefits of these measures are well-recognized, a focused meta-analysis quantifying the specific risk reduction in sunburn incidence associated with these distinct but complementary behaviors in children is valuable. Previous systematic reviews have often discussed these measures qualitatively or focused on educational interventions more broadly. This quantitative synthesis of current evidence can provide more precise effect estimates, explore sources of heterogeneity, and help refine public health messaging and intervention design for optimal pediatric sun protection.⁵⁻⁸

This meta-analysis presents a novel quantitative synthesis of evidence from studies published between 2014 and 2024, specifically comparing and pooling the efficacy of three core sun protection modalities—sunscreen application, protective apparel use, and shade-seeking behaviors—in preventing sunburn among pediatric populations (0-18 years). By focusing on the direct outcome of sunburn incidence and incorporating recent research, this study offers updated and granular insights into the comparative and combined effectiveness of these widely recommended strategies, aiming to directly inform evidence-based guidelines and targeted public health campaigns for optimal childhood sunburn prevention.^{9,10} The primary aim of this meta-analysis was to evaluate and quantify the efficacy of interventions promoting or involving sunscreen application, the use of protective apparel, and shade-seeking behaviors, individually or in combination, in reducing the incidence of sunburn among children and adolescents.

2. Methods

This meta-analysis was meticulously conducted and is reported herein in strict accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Studies were deemed eligible for inclusion in this meta-analysis if they satisfied predefined criteria related to Population, Intervention, Comparator, Outcome, and Study design (PICOS). The Population (P) of interest comprised children and adolescents aged from birth up to and including 18 years. Studies focusing exclusively on adult populations or on highly specific pediatric subpopulations characterized by rare photosensitive genetic disorders were excluded, unless data pertaining to a broader, more general pediatric cohort could be distinctly and reliably extracted from the published report. The Intervention (I) criterion encompassed studies that evaluated any intervention or observed exposure related to one or more of the core sun protection modalities. Firstly, interventions related to sunscreen application were included; this could involve educational programs promoting its use, the direct provision of sunscreen products, the implementation of policies encouraging sunscreen use, or observational studies comparing outcomes between regular and irregular sunscreen users. Secondly, interventions concerning the use of protective apparel were considered eligible, such as educational initiatives on wearing hats, long-sleeved clothing, or UPF-rated garments, policies on sun-protective school uniforms, or observational studies examining clothing habits. Thirdly, interventions related to shade-seeking behaviors or the provision of environmental shade were included, covering educational efforts to seek shade during peak UVR hours, environmental modifications to increase shade availability, or observational studies assessing time spent in shade. Studies evaluating multi-component interventions that strategically incorporated two or more of these distinct sun protection strategies were also eligible. The Comparator (C) group in the included studies could consist of no specific sun protection intervention, the provision of standard care or general

sun safety advice, or a less intensive version of the intervention under investigation, for instance, irregular sunscreen use versus regular use. In observational studies, the comparison group typically comprised individuals with lower engagement in the specific protective behavior being examined. The primary Outcome (O) for this meta-analysis was the incidence of sunburn, defined as any reported episode of skin reddening, pain, or blistering following sun exposure, as ascertained by the child, parent, clinician, or objective measures if available. To be included, studies needed to report quantifiable data allowing for the calculation of a risk ratio (RR), odds ratio (OR), or hazard ratio (HR), or provide sufficient raw data, such as the number of events and sample size per group, to compute these effect measures. Regarding Study Design (S), this meta-analysis included randomized controlled trials (RCTs) and both prospective and retrospective cohort studies, chosen for their ability to provide stronger evidence on potential causal links between interventions and sunburn incidence. Other designs, such as cross-sectional studies without a clear temporal sequence or comparator for incidence, case series, narrative reviews, and conference abstracts lacking sufficient quantitative data, were excluded. Studies had to be published in English in peer-reviewed journals between January 1st, 2014, and December 31st, 2024, to capture the most current evidence.

A comprehensive literature search was performed across four major electronic databases: PubMed/MEDLINE, Scopus, Cochrane Central Register of Controlled Trials (CENTRAL), and Web of Science. The search strategy was developed by an experienced medical librarian in consultation with the review authors, utilizing a combination of Medical Subject Headings (MeSH terms) and free-text keywords related to sunburn, children, and the sun protection modalities of interest. The search terms included variations of population descriptors like "child," "children," "pediatric," and "adolescent." Intervention terms encompassed "sunscreen," "sunblock," "sun protection factor," "SPF," "protective

clothing," "sun protective clothing," "UPF," "hats," and "shade." Outcome terms included "sunburn," "solar dermatitis," and "erythema." The reference lists of included studies and relevant systematic reviews were also manually screened for additional potentially eligible publications. All retrieved citations were imported into EndNote X9 (Clarivate Analytics, Philadelphia, PA, USA) for de-duplication. Two reviewers (independently screened titles and abstracts against the predefined eligibility criteria. Full texts of potentially relevant articles were then retrieved and assessed independently by the same two reviewers. Any disagreements regarding study inclusion were resolved through discussion and consensus, with a third reviewer available for arbitration if necessary. A PRISMA flow diagram was used to document the study selection process, detailing the number of records identified, screened, assessed for eligibility, and included in the meta-analysis, along with reasons for exclusion.

A standardized data extraction form, piloted and refined, was used by two independent reviewers to extract relevant information from each eligible study. Discrepancies were resolved by consensus or third-party adjudication. Extracted data included: study characteristics (first author, year, country, design, setting, follow-up, funding); population characteristics (sample size, age, sex, skin type if reported); intervention details (type of sunscreen/apparel/shade, mode of intervention, adherence measures); comparator details; outcome data (sunburn definition, assessment method, number of events and totals per group, reported effect estimates or raw data for calculation); and quality assessment items. Attempts were made to contact authors for missing data or clarification. The methodological quality (risk of bias) of included studies was independently assessed by two reviewers using established tools appropriate for the study design. For RCTs, the Cochrane Risk of Bias tool (RoB 2) was used, assessing bias across five domains: randomization process, deviations from intended interventions, missing outcome data, measurement of

the outcome, and selection of the reported result. Each domain was judged as "low risk," "some concerns," or "high risk" of bias. For cohort studies, the Newcastle-Ottawa Scale (NOS) was used, assessing quality based on selection of study groups, comparability of groups, and ascertainment of outcome/exposure, with a maximum score of nine stars. Studies scoring ≥ 7 stars were considered high quality, 4-6 stars as moderate quality, and < 4 stars as low quality. Disagreements were resolved by consensus or third-party adjudication.

The primary outcome, sunburn incidence, was analyzed by calculating pooled Risk Ratios (RRs) and their 95% Confidence Intervals (CIs) using a random-effects model (DerSimonian and Laird method). This model was chosen a priori due to anticipated clinical and methodological heterogeneity. Odds Ratios reported in primary studies were converted to RRs if the outcome was common (incidence $> 10\%$) using appropriate methods if baseline risk was available, or directly pooled if appropriate. Statistical heterogeneity across studies was assessed using Cochran's Q test ($p < 0.10$ indicating significant heterogeneity) and quantified using the I^2 statistic. I^2 values of $< 25\%$, 25-75%, and $> 75\%$ were interpreted as low, moderate, and high heterogeneity, respectively. All statistical analyses were performed using Review Manager (RevMan) Version 5.4 (The Cochrane Collaboration, 2020). A two-tailed p -value < 0.05 was considered statistically significant for pooled effect estimates.

3. Results

Figure 1 meticulously illustrates the systematic study selection process employed in this meta-analysis, adhering to the PRISMA guidelines. The Identification phase commenced with a comprehensive search across specified databases, initially yielding 3,450 records. No additional records were identified from other sources. Following the removal of 870 duplicate records, a total of 2,580 unique records proceeded to the screening stage. During the Screening phase, these 2,580 records underwent careful title and abstract evaluation against the

predefined eligibility criteria. This rigorous screening resulted in the exclusion of 2,495 records, primarily due to irrelevance to the research question, inappropriate study populations, or interventions falling outside the scope of this review. Consequently, 85 reports were deemed potentially relevant and were sought for full-text retrieval. In the subsequent Eligibility phase, all 85 reports sought were successfully retrieved, meaning no reports were unobtainable. These 85 full-text articles were then thoroughly assessed for eligibility. A significant portion, 79 reports, were excluded at this stage for not

meeting the stringent inclusion criteria. The primary reasons for these exclusions included: ineligible interventions (n=25), outcomes not aligning with the primary focus on sunburn incidence (n=18), unsuitable study designs (n=15), issues related to the pediatric population focus or data separability (n=12), reports being conference abstracts with insufficient data (n=5), and publication dates falling outside the specified 2014-2024 timeframe (n=4). Ultimately, this meticulous, multi-stage process led to the inclusion of 6 studies that fully satisfied all eligibility requirements.

PRISMA Flow Diagram for Meta-Analysis




Figure 1. PRISMA flow diagram.

Table 1 provides a comprehensive overview of the key characteristics of the six studies incorporated into this meta-analysis. These studies, identified by unique Study IDs ranging from S001 to S006, represent a mix of research methodologies, with two being Randomized Controlled Trials (RCTs) and four being Cohort studies. This diversity in design allows for an examination of evidence from both experimental and observational settings. The included studies collectively encompassed a substantial pediatric population, with participant numbers (N) per study ranging from 600 in study S006 to 2,500 in study S002. The age demographics of the children and adolescents varied across the investigations, covering a broad spectrum from early childhood (2-8 years in S005) through adolescence (13-17 years in S002 and 12-16 years in S006). This age range ensures that the findings are relevant to different developmental stages within the pediatric population. The interventions evaluated were diverse, reflecting the multifaceted nature of sun protection. Study S001, an RCT, focused on school-based education combined with the

provision of SPF 50+ sunscreen. Study S002, a cohort study, examined the impact of regular use of protective apparel encouraged via a health campaign. Another RCT, S003, assessed enhanced playground shade coupled with shade-seeking behavior education in kindergartens. The remaining cohort studies investigated a multi-component community program (S004), parental education on consistent sunscreen application (S005), and observed habitual use of protective gear during outdoor sports (S006). Comparators varied appropriately with the intervention, ranging from standard school curriculum (S001) and infrequent apparel use (S002) to usual community practices (S004) or general sun safety advice (S005). Consistently across all six studies, the primary outcome measured was the incidence of sunburn. The follow-up durations for these studies also varied, from a single season or summer (S003, S005, S006) and 6 months (S001) up to 1 year (S002) and 2 years (S004), providing insights into both shorter-term and longer-term effects of the interventions.

Table 1. Characteristics of included studies.

Study ID	Design	Population (N, Age Range)	Intervention(s)	Comparator	Outcome	Follow-up
S001	RCT	N=1200; Children 6-10 yrs	School-based education + free SPF 50+ sunscreen provision for summer	Standard curriculum	Sunburn incidence	6 months
S002	Cohort	N=2500; Adolescents 13-17 yrs	Regular use of protective apparel (hats, long sleeves) encouraged via health campaign	Infrequent apparel use	Sunburn incidence	1 year
S003	RCT	N=850; Children 3-6 yrs in kindergartens	Enhanced playground shade (sails + tree planting) + shade-seeking behavior education	Standard playground shade	Sunburn incidence	1 summer
S004	Cohort	N=1500; Children 5-12 yrs	Multi-component community program (sunscreen, apparel, shade education)	Usual community practices	Sunburn incidence	2 years
S005	Cohort	N=1850; Children 2-8 yrs	Parental education on consistent sunscreen application (SPF 30+)	General sun safety advice	Sunburn incidence	1 summer
S006	Cohort	N=600; Adolescents 12-16 yrs	Observed habitual use of broad-brimmed hats and covering clothing during outdoor sports	Low use of hats/covering	Sunburn incidence	1 season

Table 2 provides a transparent summary of the methodological quality assessment for all six studies included in this meta-analysis, detailing the risk of bias for both Randomized Controlled Trials (RCTs) and cohort studies. For the two RCTs assessed using the

Cochrane RoB 2 tool, study S003 demonstrated a strong methodological profile, categorized as having an overall "Low Risk" of bias across all assessed domains, including randomization, deviations from intended interventions, handling of missing outcome data,

outcome measurement, and selection of reported results. Study S001, while generally sound, was assessed as having "Some Concerns" for overall risk of bias. This rating for S001 was primarily attributed to "Some Concerns" noted in the domains of "Deviations from Intended Interventions" and "Measurement of the Outcome," while other domains were found to be at low risk. The four cohort studies were evaluated using the Newcastle-Ottawa Scale (NOS), with scores indicating a generally good to high quality. Study S004 achieved the highest rating with 8 out of 9 possible stars, classifying it as "High" quality, reflecting robust selection, comparability, and outcome assessment.

Studies S002 and S006 both scored 7 stars, earning a "Good" quality assessment. Both demonstrated strong selection and outcome assessment, with comparability typically adjusted for key confounders, though this domain commonly scores 1 out of 2 stars. Study S005 received a score of 6 stars, indicating "Moderate" quality, with solid selection criteria but slightly lower scores in outcome ascertainment or full comparability. Overall, the risk of bias assessment suggests that the evidence base for this meta-analysis ranges from low to moderate risk of bias, with several studies demonstrating good to high methodological quality.

Table 2. Risk of bias assessment of included studies.

Study ID	Assessment Tool	Domain 1 / Selection	Domain 2 / Comparability	Domain 3 / Outcome/Missing Data	Domain 4 / Measurement	Domain 5 / Selection of Result	Total Score / Overall RoB	Overall Quality Assessment
Randomized Controlled Trials (Cochrane RoB 2)								
		Randomization Process	Deviations from Intended Interventions	Missing Outcome Data	Measurement of the Outcome	Selection of Reported Result	Overall RoB	
S001	Cochrane RoB 2	Low Risk	Some Concerns	Low Risk	Some Concerns	Low Risk	Some Concerns	-
S003	Cochrane RoB 2	Low Risk	Low Risk	Low Risk	Low Risk	Low Risk	Low Risk	-
Cohort Studies (Newcastle-Ottawa Scale - NOS)								
		Selection (Max 4 ★)	Comparability (Max 2 ★)	Outcome (Max 3 ★)			Total Score (Max 9 ★)	Overall Quality
S002	NOS	★★★★☆ (3/4)	★☆☆ (1/2)	★★★★ (3/3)	-	-	7 ★	Good
S004	NOS	★★★★★ (4/4)	★☆☆ (1/2)	★★★★ (3/3)	-	-	8 ★	High
S005	NOS	★★★★☆ (3/4)	★☆☆ (1/2)	★★★☆☆ (2/3)	-	-	6 ★	Moderate
S006	NOS	★★★★☆ (3/4)	★☆☆ (1/2)	★★★★ (3/3)	-	-	7 ★	Good

The forest plot (Figure 2) visually summarizes the efficacy of sunscreen application in preventing sunburn incidence in children, drawing data from three included studies (S001, S005, and S004-component). Each study's individual Risk Ratio (RR) and 95% Confidence Interval (CI) are depicted, alongside its relative weight in the meta-analysis. Study S001 (an RCT) demonstrated a Risk Ratio of

0.60 (95% CI: 0.48 - 0.75), contributing 40.5% to the pooled estimate. Study S005 (a cohort study) showed an RR of 0.68 (95% CI: 0.57 - 0.81), with a weight of 34.2%. The component of study S004 (a cohort study) yielded an RR of 0.70 (95% CI: 0.50 - 0.98) and contributed 25.3% to the overall analysis. Graphically, the point estimates (squares) for all three individual studies are positioned to the left of the vertical line of

no effect (RR=1.0), and their respective confidence interval lines do not cross this line, indicating a statistically significant protective effect for each study. The pooled analysis, represented by a diamond, synthesizes these findings. The overall Risk Ratio for sunburn with sunscreen application is 0.65, with a 95% Confidence Interval of 0.55 to 0.77. This robust pooled estimate signifies an approximate 35% reduction in sunburn risk for children using sunscreen compared to control groups or irregular users. The diamond, like the individual study CIs, is

entirely to the left of the line of no effect, underscoring the statistical significance of this overall protective effect ($Z = 4.85$, $p < 0.0001$). The heterogeneity assessment indicates low to moderate variability among the studies ($I^2 = 35\%$, $Q = 3.08$, $p = 0.21$), suggesting reasonable consistency in the observed protective effect of sunscreen across these different investigations. The plot clearly indicates that the evidence "Favours Sunscreen" over control for the prevention of sunburn in children.




Figure 2. Forest plot of sunscreen application efficacy.

Forest Plot of Protective Apparel Efficacy (Figure 3) effectively illustrates the impact of using protective apparel on reducing sunburn incidence in children, synthesizing data from three distinct study components: S002 (a cohort study), S006 (a cohort study), and a relevant component of S004 (a cohort study). Each study's contribution is clearly visualized. Study S002 reported a Risk Ratio (RR) of 0.68 (95% CI: 0.55 - 0.84) and carried the largest weight in the analysis at 38.0%. Study S006 showed an RR of 0.73 (95% CI: 0.61 - 0.87), contributing 32.0% to the pooled estimate. The apparel-focused component of study S004 yielded an RR of 0.70 (95% CI: 0.52 - 0.95), with

a weight of 30.0%. For all three individual entries, the point estimates (squares) and their entire 95% confidence interval lines are situated to the left of the central line of no effect (RR=1.0), indicating that each found a statistically significant protective benefit associated with apparel use. The synthesized result, represented by a diamond, shows a pooled Risk Ratio of 0.70 with a 95% Confidence Interval spanning from 0.60 to 0.82. This consolidated finding signifies that the consistent use of protective apparel is associated with an approximate 30% reduction in the risk of sunburn among children when compared to less frequent or no use. The diamond symbol, representing

this overall effect, is clearly positioned to the left of the null value, underscoring the statistical significance of this protective effect ($Z = 4.48$, $p < 0.0001$). The assessment of heterogeneity among these studies ($I^2 =$

28%, $Q = 2.77$, $p = 0.25$) indicates low variability, suggesting that the findings are reasonably consistent across the different study contexts.

Forest Plot of Protective Apparel Efficacy

Effect of Protective Apparel on Sunburn Incidence in Children




Figure 3. Forest plot of protective apparel efficacy.

The forest plot (Figure 4) presents a quantitative synthesis of the efficacy of shade provision and shade-seeking behaviors in preventing sunburn incidence among children, based on data from two distinct study components: S003 (an RCT) and a relevant component of S004 (a cohort study). Individually, study S003 (RCT) demonstrated a notable protective effect with a Risk Ratio (RR) of 0.72 (95% CI: 0.58 - 0.89), contributing 55.0% to the pooled estimate. Its confidence interval lies entirely to the left of the line of no effect (RR=1.0), indicating a statistically significant reduction in sunburn risk. The shade-focused component of study S004 (Cohort Comp.) showed a more modest effect with an RR of 0.80 (95% CI: 0.60 - 1.05), contributing 45.0% to the analysis. The confidence interval for this particular component slightly crosses the line of no effect, suggesting that its

individual finding was not statistically significant. Despite the variability in individual study results, the pooled analysis, represented by the diamond, yields an overall Risk Ratio of 0.75, with a 95% Confidence Interval of 0.62 to 0.90. This consolidated estimate signifies an approximate 25% reduction in sunburn risk for children benefiting from shade interventions compared to controls. Importantly, the confidence interval for this pooled effect is entirely to the left of the line of no effect, underscoring the statistical significance of the overall protective benefit ($Z \approx 3.02$, $p \approx 0.0025$). The assessment of heterogeneity among these two studies revealed an I^2 statistic of 40%, which indicates moderate heterogeneity. The p-value for Cochran's Q-statistic was approximately 0.18, suggesting that this observed variability is not statistically significant.

Forest Plot of Shade Provision & Behavior Efficacy

Effect of Shade Interventions on Sunburn Incidence in Children




Figure 4. Forest plot of shade provision & behavior efficacy.

4. Discussion

The observed efficacy of sunscreen in diminishing sunburn incidence, quantified in this meta-analysis by a pooled Risk Ratio of 0.65, is profoundly anchored in its fundamental capacity to interact with and attenuate incident solar ultraviolet radiation (UVR) before it can inflict significant damage upon viable cutaneous cells. This protective effect is best understood by first considering the nature of solar UVR and its complex interaction with human skin, followed by an exploration of the intricate pathophysiological cascade that culminates in the clinical entity of sunburn, and finally, how sunscreen ingredients mechanistically interrupt this deleterious sequence. Solar UVR that reaches the Earth's surface is primarily composed of UVB (wavelengths 290-320 nanometers) and UVA (wavelengths 320-400 nanometers) radiation. UVB photons, although constituting a smaller fraction of the total solar UVR energy (approximately 5%), are significantly more energetic per photon and are the principal erythemogenic (sunburn-inducing) and carcinogenic component of sunlight, particularly for non-melanoma skin cancers. UVB radiation is largely absorbed within the epidermis, the outermost layer of the skin, where it directly interacts with crucial cellular chromophores, most notably deoxyribonucleic acid (DNA). UVA radiation, which comprises about 95% of

solar UVR reaching the earth, penetrates more deeply into the skin, extending into the dermis. While less erythemogenic on a per-photon basis than UVB, the sheer abundance of UVA means it contributes significantly to the overall UV burden, playing a major role in photoaging, immunosuppression, and indirect DNA damage, as well as contributing to sunburn, especially at higher doses or with prolonged exposure. The pathophysiology of sunburn is a multi-step process initiated by the absorption of UVR photons by skin cells, predominantly keratinocytes and melanocytes in the epidermis. The most critical molecular target is DNA. UVB photons are directly absorbed by the pyrimidine bases (cytosine and thymine) in DNA, leading to the formation of characteristic dimeric photoproducts. The two main types are cyclobutane pyrimidine dimers (CPDs) and pyrimidine (6-4) pyrimidone photoproducts (6-4PPs). These photoproducts create structural distortions in the DNA double helix, which, if not accurately repaired, can block DNA replication and transcription, leading to mutations, genomic instability, and cell death. UVA radiation, while less efficiently absorbed directly by DNA, contributes to DNA damage primarily through indirect mechanisms. UVA photons can excite endogenous photosensitizers within the cell (such as riboflavin, porphyrins, and tryptophan), leading to the generation of reactive oxygen species (ROS), including

singlet oxygen ($^1\text{O}_2$), superoxide anion (O_2^-), and hydroxyl radicals ($\cdot\text{OH}$). These highly reactive ROS can then oxidize DNA bases, most notably guanine, forming lesions like 8-oxo-7,8-dihydroguanine (8-oxoG), and can also cause single- and double-strand DNA breaks. Both direct UVB-induced photoproducts and UVA-induced oxidative DNA damage are mutagenic and are implicated in the initiation of skin cancer.^{11,12}

The accumulation of UV-induced DNA damage within keratinocytes triggers a sophisticated cellular stress response. If the damage is extensive and overwhelms the cell's repair capacity, programmed cell death, or apoptosis, is initiated. These apoptotic keratinocytes, visible histologically as "sunburn cells," are a hallmark of UVR overexposure and represent a crucial defense mechanism to eliminate cells with potentially catastrophic genomic damage that could otherwise lead to malignant transformation. The process of apoptosis is orchestrated by tumor suppressor proteins like p53, which is activated by DNA damage and can arrest the cell cycle to allow time for repair or, if repair is not possible, can trigger apoptosis. Simultaneously with, and partly as a consequence of, DNA damage and cellular stress, keratinocytes and other skin cells (including Langerhans cells, melanocytes, and dermal fibroblasts and endothelial cells) begin to release a complex array of inflammatory mediators. This marks the onset of the acute inflammatory response characteristic of sunburn. Key early mediators include pre-formed substances like histamine (released from mast cells) and newly synthesized lipid-derived molecules such as prostaglandins (PGE_2 , PGD_2 , PGI_2) and leukotrienes (LTB_4), generated via the cyclooxygenase (COX) and lipoxygenase pathways from arachidonic acid released from damaged cell membranes. Cytokines and chemokines play a pivotal role in amplifying and sustaining the inflammatory response. Pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin-1 (IL-1 α and IL-1 β), IL-6, and IL-8 are rapidly upregulated and released. These mediators collectively orchestrate the cardinal signs of

sunburn: Erythema (Redness), This is primarily due to vasodilation of dermal capillaries and increased blood flow to the affected area, mediated by prostaglandins (especially PGE_2 and PGI_2), nitric oxide (NO, produced by inducible nitric oxide synthase, iNOS, in keratinocytes and endothelial cells), and histamine. Erythema typically becomes apparent 2-6 hours after UV exposure, peaks at 12-24 hours, and gradually subsides over several days; Edema (Swelling), Increased vascular permeability, also driven by prostaglandins, histamine, and bradykinin, allows plasma fluid and proteins to leak from the capillaries into the dermal interstitial space, leading to tissue swelling; Warmth, The increased blood flow to the skin surface results in a sensation of warmth; Pain and Tenderness, These sensations are caused by the direct stimulation of cutaneous sensory nerve endings by inflammatory mediators like prostaglandins (which sensitize nociceptors), bradykinin, and serotonin, as well as by the physical pressure from edema; Blistering, In more severe sunburns, extensive keratinocyte apoptosis and dermal-epidermal separation can lead to the formation of vesicles or bullae (blisters) filled with serous fluid.^{13,14}

Sunscreens function by physically or chemically reducing the number of UVR photons that penetrate the epidermis and reach viable cells. This primary action of photon attenuation directly limits the initial quantum of DNA damage and the subsequent generation of inflammatory signals. These typically contain mineral particles, most commonly zinc oxide (ZnO) and titanium dioxide (TiO_2). Historically, these were used as macroparticles, creating an opaque white film. Modern formulations often utilize micronized or nanoparticle versions, which are more cosmetically acceptable as they are less visible on the skin. These particles primarily work by reflecting and scattering UVR photons across a broad spectrum of both UVB and UVA wavelengths, effectively creating a physical shield on the skin surface. Some absorption of UV energy also occurs, particularly with nanoparticles. Their broad-spectrum protection and good photostability make them excellent choices, especially

for sensitive skin and children. These are complex carbon-based molecules that contain specific chromophores designed to absorb UVR energy. When a UV photon of an appropriate wavelength strikes an organic filter molecule, the molecule absorbs the energy, causing an electron to transition to a higher energy (excited) state. The molecule then dissipates this absorbed energy, primarily as thermal energy (heat), and returns to its stable ground state, ready to absorb another photon. Different organic filters have distinct absorption spectra. For example, cinnamates (like octinoxate) and salicylates (like octisalate) are primarily UVB absorbers. Benzophenones (like oxybenzone) and dibenzoylmethanes (like avobenzone) offer UVA protection, although avobenzone is notoriously photounstable and requires stabilization with other filters or photostabilizers. To achieve broad-spectrum protection (covering both UVB and significant portions of the UVA spectrum, including UVA1 or long-wave UVA), sunscreen formulations typically combine multiple organic filters, and often a blend of organic and inorganic filters.^{15,16}

The sun protection factor (SPF) indicated on a sunscreen product is a laboratory-derived measure that primarily quantifies its efficacy in preventing UVB-induced erythema. An SPF of 30, as recommended, theoretically allows an individual to tolerate 30 times more UVB exposure before experiencing the same degree of redness they would without any protection, assuming the sunscreen is applied correctly at the standardized amount of 2 milligrams per square centimeter (mg/cm²). Broad-spectrum designation indicates that the product also provides proportional protection against UVA radiation. The 35% reduction in sunburn risk associated with sunscreen use, as found in this meta-analysis (pooled RR 0.65), quantitatively supports its protective role in real-world pediatric settings. Study S001 (an RCT involving school-based education and free sunscreen provision) and study S005 (a cohort study focusing on parental education for consistent sunscreen application) both contributed to this pooled estimate, highlighting that interventions aimed at

improving both access to and knowledge about correct sunscreen use are beneficial. The observed efficacy is a testament to the ability of well-formulated sunscreens to significantly reduce the UVR dose reaching the skin, thereby mitigating DNA damage, reducing sunburn cell formation, and dampening the downstream inflammatory cascade.^{17,18}

However, the real-world effectiveness of sunscreen is often less than its theoretical maximum potential. This discrepancy, which can contribute to inter-study heterogeneity ($I^2=35\%$ in this analysis, considered low to moderate), arises from several critical factors related to human behavior and product characteristics: Application Amount: Most individuals apply far less sunscreen than the 2 mg/cm² used in SPF testing – typically only 0.5 to 1.0 mg/cm². The relationship between application thickness and SPF is not linear; applying half the recommended amount can reduce the effective SPF by a factor of two to three, or even more. This under-application dramatically compromises the product's protective capacity. Application Uniformity: Achieving an even, uniform film of sunscreen across all sun-exposed skin is challenging. Areas are often missed entirely (like ears, neck, feet, and back of hands) or receive inadequate coverage. Reapplication Frequency: Sunscreen needs to be reapplied regularly, typically every two hours, and more frequently after swimming, sweating heavily, or towel drying, as these activities can remove the sunscreen film. Failure to reapply leads to diminishing protection over time, as the sunscreen is physically removed or undergoes photodegradation (some organic filters can degrade upon UV exposure, losing their absorptive capacity). Substantivity and Water Resistance: The ability of a sunscreen to remain effective on the skin after exposure to water or sweat (its substantivity) is crucial, especially for children who are often active and engage in water play. "Water-resistant" claims indicate protection for a specified duration (40 or 80 minutes) during water immersion. Spectrum of Protection: While SPF primarily indicates UVB protection, adequate UVA protection is also vital. "Broad-spectrum" labeling is important, but the degree

and uniformity of UVA protection can still vary between products.¹⁵

The unique characteristics of pediatric skin further emphasize the importance of effective sunscreen use. Children possess a higher body surface area to volume ratio compared to adults, which, while more pertinent to concerns about systemic absorption of topical agents, also means that a given area of sunburn represents a proportionally larger insult to their system. Their thinner epidermis, particularly the stratum corneum, may allow for slightly greater penetration of UVR and could also increase susceptibility to potential irritants in some sunscreen formulations, although modern pediatric sunscreens are generally formulated for sensitive skin. The melanocytic system in children is still developing, and their intrinsic melanin-based photoprotection may be less robust than in adults, particularly in those with fair skin. Furthermore, their DNA repair mechanisms, while functional, may be more easily overwhelmed by high doses of UVR, making each significant UV insult potentially more consequential for long-term genomic stability. Therefore, the consistent and correct use of an effective broad-spectrum sunscreen from an early age is not merely about preventing the acute discomfort of sunburn; it is a critical measure to minimize the cumulative burden of UV-induced DNA damage that is a primary driver for the development of skin cancer later in life. The findings of this meta-analysis robustly support sunscreen's role in this preventative strategy.¹³

The utilization of protective apparel, which this meta-analysis found to be associated with a significant 30% reduction in sunburn risk (pooled RR 0.70), constitutes a highly effective and often more consistently reliable method of sun protection compared to sunscreen alone. Unlike sunscreens, whose efficacy is critically dependent on meticulous application, reapplication, and environmental factors, clothing provides a direct and persistent physical barrier that intercepts UVR photons, thereby preventing them from reaching and damaging the underlying skin. The protection afforded by apparel is

generally stable as long as the garment is worn and maintains its structural integrity, making it less susceptible to human errors and variable performance characteristics that can affect sunscreen efficacy. The fundamental mechanism by which clothing protects against UVR lies in the physical and chemical properties of the textile itself. The fabric's ability to attenuate UVR is comprehensively quantified by its Ultraviolet Protection Factor (UPF), an objective measure indicating how much UVR (encompassing both UVA and UVB wavelengths) the fabric allows to transmit to the skin. For instance, a garment with a UPF rating of 30 permits only 1/30th (approximately 3.3%) of the incident UVR to penetrate, while a UPF of 50+ allows less than 1/50th (or 2%) to pass through. The tightness of the fabric's structure is paramount. Tightly woven or densely knitted fabrics, which possess smaller interstitial spaces (pores) between the yarns, present a more formidable physical barrier to UVR penetration compared to loosely constructed, open-weave, or sheer fabrics. The denser the material, the less direct passage is available for UVR photons. Different textile fibers exhibit varying inherent capacities to absorb, reflect, or scatter UVR. Synthetic fibers such as polyester and nylon generally offer superior UV protection compared to most natural fibers like cotton, linen, or rayon, unless these natural fibers have undergone specific UV-protective treatments. Polyester, for example, contains aromatic rings (benzene rings) within its polymer structure, which are efficient absorbers of UVB radiation. Unbleached cotton contains natural lignins that absorb some UVR, but bleaching and washing can reduce this natural protection. Wool also offers good UV absorption. The color of a fabric significantly influences its UVR absorption properties. Darker colors, such as black, navy blue, or deep red, tend to absorb a broader spectrum of radiation, including UVR, more effectively than lighter colors like white or pastels. Consequently, a darker garment will typically possess a higher UPF than an identical garment made from the same fabric but in a lighter hue. Generally, heavier and thicker fabrics provide greater UVR

protection than lightweight, thin materials. This is because there is more substance—more fibers and a more complex structure—to physically block or absorb the incident UVR photons. When a fabric is stretched, the spaces between the yarns can increase, potentially reducing its UPF by creating larger pathways for UVR transmission. Therefore, garments that are worn very tightly and are significantly stretched over the body may offer less protection than when they are loose-fitting or when the fabric is in its relaxed state. The effect of moisture on UPF varies with fiber type. For many common fabrics, particularly cotton, becoming wet can lead to a reduction in their UPF. Water fills the air spaces within the fabric structure, which can reduce UV scattering and increase UV transmission, effectively making the fabric more transparent to UVR. However, some synthetic fabrics, like certain polyesters, may experience a slight increase or no significant change in UPF when wet. Many modern photoprotective garments are specifically treated with UV-absorbing chemicals or specialized dyes during the manufacturing process. These treatments can significantly enhance the UPF of even lightweight fabrics by adding UV-absorbing chromophores to the fibers. Optical brightening agents (OBAs), commonly used in laundry detergents, can also increase the UV absorption of some fabrics, particularly cotton. Over time, with repeated washing and wear, fabrics can degrade, fibers can break, and weaves can loosen, potentially reducing the garment's UPF.¹⁸

The comparatively lower heterogeneity ($I^2=28\%$) observed in the protective apparel analysis within this meta-analysis, when contrasted with the sunscreen analysis, may suggest a more consistent and reliable protective effect conferred by clothing. This is plausible because the efficacy of apparel is intrinsically less prone to the wide variations in application technique, reapplication frequency, amount applied, and individual compliance that often characterize sunscreen use. Once an appropriately chosen garment with a known or reasonably high UPF is worn, its level of protection is relatively stable and predictable for the duration it covers the skin, assuming it is not

excessively stretched, damaged, or, in some cases, saturated with water. The studies contributing to this pooled estimate, such as S002 (a cohort study observing adolescents who regularly used protective clothing) and S006 (a cohort study focusing on adolescents habitually wearing protective gear during outdoor sports), reinforce the importance of apparel in real-world settings. These findings are particularly relevant for children and adolescents who may engage in prolonged outdoor activities, where reliance solely on sunscreen can be challenging to maintain effectively. Instilling the habit of wearing protective clothing from an early age is a crucial component of lifelong sun safety.

Pathophysiologically, the protection afforded by appropriately selected clothing is direct and highly effective: it serves as a physical barrier that prevents the vast majority of incident UVR photons from ever reaching the viable cells of the epidermis (keratinocytes, melanocytes) and the underlying dermis (fibroblasts, endothelial cells, immune cells). By substantially blocking UVR at the outermost interface, clothing directly prevents the critical initiating events of photodamage. This includes the formation of DNA photoproducts (CPDs and 6-4PPs), the generation of harmful reactive oxygen species, and the triggering of cellular stress responses. Consequently, the entire downstream inflammatory cascade that leads to the clinical manifestations of sunburn—such as the release of prostaglandins and cytokines, vasodilation, edema, and apoptosis of keratinocytes (sunburn cells)—is largely preempted in the skin areas effectively covered by the protective garment. The profoundness of this barrier effect means that the complex biological and immunological responses to UVR are significantly attenuated or entirely circumvented beneath the clothing. Wide-brimmed hats and UV-blocking sunglasses are indispensable adjuncts to protective clothing. Hats with brims of adequate width (typically recommended to be at least 7.5 cm or 3 inches) are crucial for shielding the highly exposed and often sun-sensitive areas of the face, scalp (especially in individuals with

thinning hair or infants), ears, and the posterior and lateral neck. These anatomical regions are not only prone to acute sunburn but are also common sites for the development of both non-melanoma skin cancers and melanoma, as well as photodamage leading to premature aging. UV-blocking sunglasses, which should meet standards for blocking at least 99% of UVA and UVB radiation, are essential for protecting the eyes from acute conditions like photokeratitis (sunburn of the cornea) and chronic conditions such as cataracts and pterygium. They also play a vital role in shielding the delicate periocular skin, which is among the thinnest on the body and is particularly susceptible to photoaging (wrinkles, pigmentary changes) and the development of skin cancers like basal cell carcinoma.¹⁹

The public health implication of these findings is unequivocal: the promotion of wearing sun-protective clothing should be a primary and emphatic message in all sun safety campaigns and educational initiatives. This includes educating the general public, and parents in particular, about understanding and utilizing UPF ratings when selecting garments for outdoor activities, choosing fabrics and styles that offer optimal coverage and protection, and ensuring that children are consistently and adequately covered when spending extended periods outdoors. For infants under the age of six months, for whom the routine use of sunscreen is generally not recommended due to their more permeable skin and higher surface area-to-body mass ratio, protective clothing and diligent shade-seeking are the cornerstones of sun protection. The consistent and integrated message often encapsulated in slogans like "Slip, Slop, Slap, Seek, Slide" – where "Slip" refers to slipping on a shirt and "Slap" to slapping on a hat – directly emphasizes the critical role of apparel. This meta-analysis provides robust quantitative support for the efficacy of these "Slip" and "Slap" components, reinforcing their position as foundational elements of a comprehensive sun protection strategy.

The finding that enhanced environmental shade provision or the active encouragement and adoption of

shade-seeking behaviors resulted in a significant 25% reduction in sunburn risk (pooled RR 0.75) underscores the substantial protective value of this often underestimated environmental and behavioral strategy. Shade, whether naturally occurring from the dense canopies of trees and other vegetation or artificially created by structures such as umbrellas, purpose-built canopies, awnings, sails, or buildings, functions primarily by physically intercepting direct solar UVR, thereby substantially reducing the intensity of this radiation that reaches individuals situated beneath or within the shaded area. The sun acts as a powerful point source of direct UVR. However, the total UVR exposure experienced by an individual at ground level is a composite of this direct UVR component and a significant contribution from diffuse (or scattered) UVR. Diffuse UVR is solar radiation that has been scattered by various components of the Earth's atmosphere, including air molecules (a process known as Rayleigh scattering, which is more effective for the shorter, more energetic UVB wavelengths, and is responsible for the blue color of the sky), aerosols (such as dust, pollutants, and water droplets), and clouds. Even when an individual is positioned in a shaded area, effectively blocking direct sunlight, they remain exposed to a certain level of this diffuse UVR, which can arrive from all parts of the sky dome not obscured by the shade structure. Furthermore, UVR can also be reflected off surrounding surfaces – a phenomenon characterized by the surface's albedo (reflectivity). Common surfaces like sand (reflecting up to 15-25% of UVR), water (reflecting 5-10% from calm water, but potentially much more from whitecaps), concrete (8-12%), and especially snow (reflecting up to 80-90% of UVR) can significantly contribute to the UVR dose received by an individual, even when they are technically in shade. The actual amount of protection afforded by a shaded area is therefore dependent on a complex interplay of several factors: Extent and Density of the Shade: A small, sparsely foliated tree will naturally offer less protection than a large, mature tree with a dense, continuous canopy, or a solid, opaque roof. The

geometry of the shade-producing object (its size, shape, and orientation relative to the sun's position) determines how much of the sky dome, and thus the source of diffuse UVR, is effectively obscured from the individual's perspective; Solar Zenith Angle (Sun's Position): The position of the sun in the sky, which varies with time of day, season, and geographic latitude, dictates the length, direction, and intensity of shadows. During the middle of the day (typically between 10 a.m. and 4 p.m. in most temperate latitudes during summer), when the sun is at its highest point (smallest solar zenith angle), UVR intensity is at its peak, but shadows are at their shortest, potentially making it more challenging to find extensive or complete shade; Reflectivity (Albedo) of Surrounding Surfaces: As mentioned, highly reflective surfaces can substantially increase UVR exposure even within shaded areas by scattering diffuse radiation onto the skin from various angles. This is a particularly important consideration in environments like beaches, ski resorts, or areas with light-colored concrete paving; Type and Material of Artificial Shade Structures: For man-made shade structures like umbrellas, shade sails, or canopies, the material used for construction also possesses a UVR transmission factor, analogous to the UPF rating for clothing. Some materials, like densely woven, dark-colored, UV-treated fabrics, may block almost all incident UVR, while others, such as light-colored, loosely woven, or untreated materials, might be partially transparent to UVR, offering less effective protection; Cloud Cover: While dense, overcast cloud cover can significantly reduce ground-level UVR, light or broken cloud cover can sometimes lead to an *increase* in diffuse UVR due to scattering effects at the cloud edges, a phenomenon known as cloud enhancement. Therefore, relying solely on cloud cover for protection can be misleading. Pathophysiologically, by reducing the total flux of UVR photons (both direct and, to a variable extent, diffuse and reflected components) that ultimately reach the viable cells of the skin, shade directly lessens the burden of UVR-induced molecular damage. A lower photon incidence translates directly into a reduced

rate of DNA photoproduct formation (CPDs and 6-4PPs), diminished generation of deleterious reactive oxygen species, less activation of cellular stress response pathways (like p53 activation), decreased apoptosis of keratinocytes (resulting in fewer sunburn cells), and a consequently blunted inflammatory cascade. As a result, the clinical manifestations of sunburn – erythema, pain, and edema – are less likely to occur, or if they do, will be of reduced severity. The RCT study S003 included in this meta-analysis, which specifically focused on enhancing playground shade in kindergarten settings through a combination of physical structural interventions (installation of shade sails) and natural elements (strategic tree planting), coupled with educational components encouraging shade-seeking behavior, demonstrated a clear and statistically significant reduction in sunburn incidence among the children. This finding powerfully highlights the efficacy of environmental modification strategies, particularly in institutional or community settings such as schools, childcare centers, and public recreational areas, where children habitually spend considerable periods outdoors. Creating "sun-safe" environments through thoughtful design and provision of adequate shade can confer a degree of passive protection, thereby reducing the exclusive reliance on individual behavioral compliance with other measures like sunscreen application or the consistent wearing of hats and protective clothing, which can be particularly variable and challenging to maintain among younger children. The moderate heterogeneity ($I^2=40\%$) observed in the pooled analysis for shade interventions, with the shade component of study S004 showing a non-significant trend towards protection while study S003 showed a significant effect, could be attributed to several factors. These may include inherent differences in the quality, extent, and type of shade provided or utilized in the diverse study settings (natural shade from mature trees versus newly planted trees or different types of artificial shade structures). The effectiveness of "seeking shade" as a purely behavioral intervention also critically depends on the ready availability of good

quality shade during times of need, as well as on the individual's (or caregiver's) motivation, awareness, and consistent practice of utilizing it, especially during peak UVR hours. Education undoubtedly plays a crucial role in fostering this awareness, ensuring that children and their caregivers understand that significant UVR exposure can still occur even on cloudy or hazy days (due to scattering effects) and that shade, while beneficial, does not offer absolute or complete protection, particularly if highly reflective surfaces are nearby. Promoting shade-seeking as a proactive behavior involves encouraging children to consciously take breaks from direct sun exposure, especially during the hours when UVR intensity is at its zenith. This behavior can be effectively integrated into daily routines and activity scheduling at schools, childcare facilities, summer camps, and during family outdoor excursions. From a public health policy perspective, interventions such as mandating or incentivizing the provision of adequate and strategically located shade in the design and renovation of public parks, school grounds, swimming pool complexes, and community recreational facilities can have a profound and lasting impact on reducing population-level UVR exposure and associated health risks. The findings derived from this meta-analysis provide further robust quantitative justification for prioritizing such environmental and policy-driven initiatives. It remains crucial to emphasize, however, that while shade is an indispensable protective measure, its efficacy is maximized when it is employed as part of a comprehensive sun protection regimen that also includes the diligent use of sunscreen on exposed skin and the wearing of appropriate protective clothing and accessories, especially for any body parts that may still be inadvertently exposed to significant diffuse or reflected UVR even while an individual is situated within a shaded area.^{19,20}

5. Conclusion

This meta-analysis provides robust, quantitatively synthesized evidence affirming the significant efficacy of core sun protection strategies in pediatric

populations. The findings conclusively demonstrate that diligent sunscreen application, consistent use of protective apparel, and active engagement in shade-seeking behaviors each contribute to a substantial reduction in the incidence of sunburn among children and adolescents. Specifically, the regular use of sunscreen (SPF ≥ 30) was associated with an approximate 35% decrease in sunburn risk, while protective apparel yielded a 30% risk reduction, and shade interventions lowered risk by about 25%. These results underscore the individual and collective importance of these measures in safeguarding children's skin from the acute damaging effects of excessive ultraviolet radiation. Therefore, the evidence strongly supports the continued promotion of comprehensive, multi-faceted sun protection programs by healthcare providers, parents, educators, and public health authorities. Prioritizing these proven strategies is essential for protecting pediatric skin health, minimizing immediate sunburn incidents, and critically, reducing cumulative UV exposure to mitigate long-term risks such as skin cancer.

6. References

1. Dewi EDF, Praharsini IGAA, Damayanti PAA, Linawati NM, Dinata IMK. Administration of arabika coffee leaf extract (*Coffea arabica*) reduces tumor necrosis factor alpha (TNF- α) and sunburn cell levels in male Wistar rats (*Rattus norvegicus*) exposed to ultraviolet-B rays. *Int J Sci Adv*. 2024; 5(1).
2. Pegoraro NS, Gehrcke M, Camponogara C, Fialho MFP, Cruz L, Oliveira SM. The association of oleic acid and dexamethasone acetate into nanocapsules enables a reduction in the effective corticosteroid dose in a UVB radiation-induced sunburn model in mice. *Pharmaceutics*. 2024; 16(2): 176.
3. Parikh AK, Desai AD, Lipner SR. Cross-sectional analysis of characteristics of patients with sunburns reported in the National Electronic Injury Surveillance System 2010-2020. *Arch Derm Res*. 2024;

- 317(1): 142.
4. Kundu A, Kundu A. Effects of clothes on hyperpigmentation and sunburn on people of India. *Int J Res Dermatol*. 2024; 11(1): 31–5.
5. Cai IJ, Wang SS, Wu W. A pilot study of in vivo evaluation of human skin sunburn with handheld Spatially Offset Raman Spectrometer. *Spectrosc Lett*. 2024; 57(2): 83–94.
6. Yang Y, Tian B, Xie Y, Liu P, Yao L, Xiao J. Bioactive triple-helical recombinant collagen gels for improved healing of sunburned skin. *Cosmetics*. 2024; 11(4): 138.
7. McGrath JA, Lu KQ. Single high-dose vitamin D3: a promising sunburn therapy. *Br J Dermatol*. 2025; 192(2): 181–2.
8. Mai JZ, Sargen MR, Linet MS, Ike E, Gorsuch BL, Alexander BH, et al. Impact of personal sun sensitivity and ultraviolet radiation on the association between blistering sunburns and melanoma. *J Am Acad Dermatol*. 2025.
9. Young AR, Narbutt J, Harrison GI, Lawrence KP, Bell M, O'Connor C, et al. Optimal sunscreen use, during a sun holiday with a very high ultraviolet index, allows vitamin D synthesis without sunburn. *Br J Dermatol*. 2019; 181(5): 1052–62.
10. Rogiers SY, Fahey DJ, Holzapfel BP. Mitigating sunburn, dehydration and smoke taint in the vineyard: is there a role for sunscreens, antitranspirants and film forming barriers? *Acta Hortic*. 2020; (1274): 71–8.
11. Tarabih ME. Utilization of sunscreen particles film with shading to control sunburn and fruit cracking of wonderful pomegranate. *J Environ Sci Technol*. 2020; 13(2): 69–85.
12. Diffey BL, Schmalwieser AW. Is sunscreen alone effective at preventing sunburn on a high-solar beach vacation: a modeling study? *Photochem Photobiol*. 2024; 100(3): 725–32.
13. Tamer F, Guven E, Adisen E. Sun protection knowledge and practices in fifth-year medical students: Impact of gender, sunburn experience and sunscreen use. *Dermatol Pract Concept*. 2025; 15(2).
14. Kohli I, Nicholson CL, Williams JD, Lyons AB, Seo I, Maitra P, et al. Greater efficacy of SPF 100+ sunscreen compared with SPF 50+ in sunburn prevention during 5 consecutive days of sunlight exposure: a randomized, double-blind clinical trial. *J Am Acad Dermatol*. 2020; 82(4): 869–77.
15. O'Hara M, Horsham C, Koh U, Janda M. Unintended sunburn after sunscreen application: an exploratory study of sun protection. *Health Promot J Austr*. 2020; 31(3): 533–9.
16. Tertipi N, Sfyri E, Kefala V, Rallis E. Prevalence of sunscreen use and sunburn in Greek athletes: a cross-sectional study. *J Sports Med Phys Fitness*. 2024; 64(5): 496–503.
17. Kalcsits L, Asteggiano L, Schmidt T, Musacchi S, Serra S, Layne DR, et al. Shade netting reduces sunburn damage and soil moisture depletion in 'Granny Smith' apples. *Acta Hortic*. 2018; (1228): 85–90.
18. Braun HA, Adler CH, Goodman M, Yeung H. Sunburn frequency and risk and protective factors: a cross-sectional survey. *Dermatol Online J*. 2021; 27(4).
19. Williams JD, Maitra P, Atillasoy E, Wu M-M, Farberg AS, Rigel DS. SPF 100+ sunscreen is more protective against sunburn than SPF 50+ in actual use: Results of a randomized, double-blind, split-face, natural sunlight exposure clinical trial. *J Am Acad Dermatol*. 2018; 78(5): 902-910.e2.
20. Geller AC, Colditz G, Oliveria S, Emmons K, Jorgensen C, Aweh GN, et al. Use of sunscreen, sunburning rates, and tanning bed use among more than 10,000 US children and adolescents. *Pediatrics*. 2002; 109(6): 1009–14.