



Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: www.bioscmed.com

The Hidden Hazard of Celebration: A Meta-Analysis of Ocular Morbidity from Colored Powder Festivals and Implications for Global Travel Medicine

Frangklin Refelino Jotlely^{1*}, I Made Ady Wirawan², I Gusti Ayu Made Juliari¹, Ida Ayu Ary Pramita¹

¹Department of Ophthalmology, Faculty of Medicine, Universitas Udayana/Prof. Dr. I.G.N.G. Ngoerah General Hospital, Denpasar, Indonesia

²Faculty of Medicine, Universitas Udayana, Denpasar, Indonesia

ARTICLE INFO

Keywords:

Chemical keratitis
Holi
Ocular trauma
Public health
Travel medicine

*Corresponding author:

Frangklin Refelino Jotlely

E-mail address:

frangklin.jotlely@gmail.com

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v9i10.1417>

ABSTRACT

Background: The globalization of colored powder festivals, such as Holi, has exposed millions, including international tourists, to significant ocular health risks from synthetic industrial dyes that have replaced traditional organic powders. This study was conducted to quantify the prevalence of acute ocular morbidities from these festivals, providing an evidence base to inform clinical practice and public health policy, particularly within travel medicine. **Methods:** Following PRISMA guidelines, a systematic search was conducted in PubMed, Google Scholar, ScienceDirect, and ProQuest for observational studies published up to July 2024. Studies reporting ocular complications from colored powder exposure were included. Data were extracted by two independent reviewers, and methodological quality was assessed using JBI checklists. A random-effects model was used to calculate pooled prevalence for key ocular injuries. Heterogeneity was investigated using sensitivity analyses, and publication bias was assessed with funnel plots. **Results:** Six studies from India, encompassing 189 patients, met the inclusion criteria for quantitative analysis. The patient population was predominantly young males (mean age 20-30 years). The meta-analysis revealed a high pooled prevalence of key ocular morbidities. The prevalence for chemical conjunctivitis was 95% (95% CI: 87%-100%), though with substantial but anticipated heterogeneity ($I^2=82\%$). The prevalence for corneal epithelial defects was 37% (95% CI: 23%-53%; $I^2=45\%$), and for superficial punctate keratopathy was 75% (95% CI: 57%-90%; $I^2=0\%$). Most injuries were bilateral. **Conclusion:** Participation in colored powder festivals, based on extensive evidence from Holi in India, presents a quantifiable and significant risk of acute ocular morbidity. The high prevalence of chemical-induced ocular surface disease necessitates urgent recognition of this "hidden hazard." Proactive, evidence-based preventive strategies, including tiered recommendations for protective eyewear and updated pre-travel health advisories, are imperative to safeguard the vision of millions of celebrants worldwide.

1. Introduction

Cultural celebrations involving the throwing of colored powders have transcended their traditional origins to become a worldwide phenomenon. The most renowned of these is the Hindu festival of Holi, the "Festival of Colours," which celebrates the arrival of spring and the triumph of good over evil.¹ Originally a religious and cultural event localized to the Indian

subcontinent, its vibrant and immersive nature has captivated a global audience. In recent decades, this practice has been adopted and commercialized into secular events such as "Color Runs" and music festivals across North America, Europe, Asia, and beyond, drawing millions of participants annually.² This globalization has transformed these events into major tourist attractions. International travelers

actively seek out these experiences, integrating them into their itineraries to engage with local culture in a deeply personal and memorable way. From an ocular travel medicine perspective, this trend represents a unique and profoundly understudied risk factor. While travel medicine guidelines provide extensive advice on infectious diseases, vaccinations, and general safety, specific warnings regarding ocular hazards at mass cultural gatherings like colored powder festivals are conspicuously absent. This oversight is critical, as tourists are often ill-prepared for the potential dangers and may face significant barriers, such as language and unfamiliarity with local healthcare systems, in seeking timely and appropriate ophthalmic care, thereby exacerbating the severity of potential injuries.³

The ocular risks associated with these festivals have escalated dramatically with a fundamental shift in the composition of the colored powders. Traditionally, Holi powders, or gulal, were prepared from safe, natural sources such as dried flowers, herbs like turmeric and neem, and other plant-based materials. These preparations were largely benign and associated with minimal ocular irritation. However, the pressures of commercialization and mass production have led to the widespread replacement of these natural ingredients with cheaper, synthetically produced industrial dyes and fillers. Chromatographic studies have confirmed the presence of a cocktail of hazardous substances in commercially available powders.⁴ These include known carcinogens and potent irritants such as Malachite Green (a textile dye), Gentian Violet, Rhodamine, and Auramine O. To add bulk and texture, manufacturers often include abrasive particulate matter like silica dust, powdered glass (mica), and asbestos. Furthermore, heavy metals, including lead oxide, mercury sulfite, and chromium iodide, have been identified as frequent adulterants, adding another layer of toxicity. The regulation of these powders is inconsistent globally. In South Asia, where Holi is most widely celebrated, powders often lack any standards of purity or safety labeling. While products for commercial events like

Color Runs in Western countries are often marketed as "certified non-toxic," this certification can be manufacturer-dependent, with incomplete oversight regarding factors like alkaline pH levels or contamination with endotoxins from bacteria. Tourists, unaware of these regional regulatory disparities, may participate with a false sense of security, failing to take necessary precautions.

Exposure of the ocular surface to these modern synthetic powders initiates a multi-faceted pathophysiological cascade, combining chemical, mechanical, and biological insults that can lead to both acute injury and chronic, debilitating sequelae. The powders are often markedly alkaline, with a pH well above the neutral 7.4 of the ocular surface. This alkalinity saponifies the fatty acids in the cell membranes of the corneal and conjunctival epithelium, a process known as liquefactive necrosis. This rapid dissolution of the cell membrane disrupts cellular integrity and allows the toxic aniline dyes and heavy metal adulterants to penetrate deep into the corneal stroma. This mechanism is the hallmark of alkali chemical burns and is what makes them notoriously more damaging than acid burns, which tend to cause coagulative necrosis that self-limits deeper penetration.⁵ The immediate response to this chemical assault is a massive inflammatory cascade. Mast cells in the conjunctiva degranulate, releasing histamine and other vasoactive mediators, leading to profound vasodilation (redness) and increased vascular permeability (chemosis). This process is further amplified by the influx of polymorphonuclear leukocytes (PMNs), a response driven by chemokines like Interleukin-8 (IL-8). These activated PMNs release proteolytic enzymes, such as matrix metalloproteinases (MMPs), and pro-inflammatory cytokines, including tumor necrosis factor-alpha (TNF- α) and Interleukin-1 beta (IL-1 β), which together orchestrate a self-perpetuating cycle of tissue degradation and inflammation.⁶ The synthetic dyes themselves, particularly substances like malachite green, are directly cytotoxic. They generate a surge of reactive oxygen species (ROS), such as superoxide

radicals and hydrogen peroxide. This "oxidative burst" overwhelms the natural antioxidant defenses of the corneal epithelium (like superoxide dismutase and catalase), leading to widespread cellular damage through lipid peroxidation of cell membranes, mitochondrial dysfunction, and DNA damage. This oxidative stress is a potent trigger for apoptosis, or programmed cell death, leading to the loss of viable epithelial cells and compromising the integrity of the ocular surface.⁷ The inclusion of particulate matter such as silica, mica, and fine glass particles acts as a mechanical abrasive. During the rubbing and throwing of colors, these sharp, hard particles scrape against the delicate ocular surface. With every blink, the eyelid drags these abrasives across the globe, physically scouring away epithelial cells. This creates myriad micro-traumas, ranging from fine punctate erosions to large, confluent epithelial defects. These defects not only cause significant pain and blurred vision but also critically break down the primary physical barrier of the eye. The powders are often produced and stored in non-sterile conditions, leading to contamination with bacteria and fungi.⁸ The presence of endotoxins, which are lipopolysaccharide components of Gram-negative bacteria's cell walls, has been confirmed in commercial powder samples. These endotoxins are potent activators of the innate immune response via Toll-like receptor 4 (TLR4), further amplifying the inflammatory cascade. When introduced into an eye with a compromised epithelial barrier from chemical and mechanical trauma, these contaminating pathogens can lead to secondary infectious keratitis and corneal ulcers, a potentially blinding complication.⁹

The existing body of literature on ocular complications from colored powder festivals consists primarily of single-center, retrospective case series and individual case reports. While these studies provide valuable descriptive insights, the findings remain fragmented and potentially heterogeneous due to differences in powder composition, patient populations, and clinical reporting standards. This fragmentation has hindered the establishment of a

clear, evidence-based understanding of the true burden of ocular morbidity. A systematic review and meta-analysis are necessary to synthesize this disparate evidence, providing a quantitative estimate of the risks involved. By pooling data across studies, we can derive more robust and generalizable estimates of the prevalence of specific injuries, identify the most vulnerable demographics, and create a solid evidence base to inform clinical practice and public health policy. The novelty of this study is threefold. First, it is the first meta-analysis to quantitatively synthesize the prevalence of specific ocular morbidities arising from colored powder festival participation, transforming descriptive accounts into statistical evidence. Second, it moves beyond a purely clinical focus by explicitly framing the issue as a global travel medicine concern, aiming to bridge the gap between clinical ophthalmology and pre-travel health advisory services. Third, by incorporating a detailed discussion of long-term sequelae and pragmatic, tiered public health recommendations, it provides a more holistic and actionable perspective than previous literature.¹⁰ The primary aim of this study was to conduct a systematic review and meta-analysis to determine the pooled prevalence of the most common acute ocular injuries—specifically chemical conjunctivitis, corneal epithelial defects, and superficial punctate keratopathy—resulting from exposure to colored powders during festivals. Secondary aims were to synthesize the demographic characteristics of the affected population, to critically evaluate the implications of these findings for the development of evidence-based guidelines for international travelers and public health organizations, and to contextualize the acute injuries within the broader spectrum of potential long-term ocular surface disease.

2. Methods

This systematic review and meta-analysis were conducted and reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. The study protocol was designed a priori to ensure

methodological rigor and transparency in all stages of the review process. A comprehensive and systematic literature search was performed across four electronic databases: PubMed, Google Scholar, ScienceDirect, and ProQuest. The search was designed to be highly sensitive to identify all relevant publications, with no date restrictions applied, encompassing all articles published up to July 2024. The search strategy combined Medical Subject Headings (MeSH) terms and free-text keywords using Boolean operators ("AND," "OR"). The core keywords included: ("Holi" OR "Holi festival" OR "colour festival" OR "color festival" OR "Colour Run" OR "colored powder") AND ("ocular" OR "eye" OR "ophthalmic") AND ("injury" OR "trauma" OR "complication" OR "keratitis" OR "conjunctivitis" OR "morbidity"). The search syntax was adapted for each database's specific indexing system to maximize retrieval. Additionally, the reference lists of all included articles and relevant narrative reviews were manually screened to identify any further studies that may have been missed in the initial electronic search.

Studies were selected for inclusion based on a predefined set of eligibility criteria, structured around the Population, Exposure, Outcome, and Study Design (PEOS) framework. Population (P): The study included individuals of any age or gender who presented to a medical facility or were examined for ocular issues temporally associated with their participation in a colored powder-based festival. Exposure (E): The exposure of interest was direct ocular contact with synthetic or natural colored powders during participation in a celebratory event like the Holi festival or a commercial Color Run. Studies focusing on injuries from other festival-related activities, such as firecrackers or explosives, were excluded. Outcome (O): The primary outcomes for meta-analysis were the prevalence of specific ocular diagnoses. For the purpose of this review, these were operationally defined as follows: Chemical Conjunctivitis: Considered present if the primary study reported the diagnosis itself, or reported the presence of conjunctival hyperemia (redness) and/or chemosis (swelling) on examination; Corneal

Epithelial Defect: Considered present if the study reported a diagnosis of corneal abrasion, corneal erosion, or a localized epithelial defect visualized with fluorescein staining; Superficial Punctate Keratopathy (SPK): Considered present if the study reported this specific diagnosis or described a diffuse, punctate pattern of fluorescein staining on the cornea. Study Design (S): Only observational studies providing prevalence data were included, such as cross-sectional studies and case series (defined as ≥ 2 patients). Case reports ($n=1$) were considered for the qualitative synthesis but were explicitly excluded from the quantitative meta-analysis due to their inability to provide a valid prevalence estimate. Excluded study types were purely experimental laboratory or animal studies, narrative reviews, systematic reviews, editorials, commentaries, and letters without primary data. Studies had to be published in English or Bahasa Indonesia to be included.

The study selection process was performed by two independent reviewers. Initially, all retrieved records were imported into a reference management software, and duplicate records were removed. The reviewers then independently screened the titles and abstracts of the remaining unique records against the eligibility criteria. Full-text articles of all potentially relevant records were subsequently retrieved and assessed for final inclusion. Any discrepancies or disagreements between the reviewers at any stage of the screening process were resolved through discussion and consensus, with a third reviewer available for arbitration if needed. A standardized data extraction form was used to systematically collect relevant information from each included study. The following data points were extracted: first author, year of publication, country of study, study design, study period, total sample size, patient demographic characteristics, and detailed clinical outcomes. For the meta-analysis, the number of patients experiencing a specific ocular morbidity (events) and the total number of patients in the study (total) were extracted for each relevant outcome. The methodological quality and risk of bias of each

included study were independently assessed by two reviewers using the appropriate Joanna Briggs Institute (JBI) Critical Appraisal Checklist corresponding to the study design. Each item on the checklists was answered with "Yes," "No," "Unclear," or "Not Applicable." Based on the proportion of "Yes" scores, studies were qualitatively categorized as having a low, moderate, or high risk of bias. This assessment was not used to exclude studies but rather to contextualize the findings and inform the investigation of heterogeneity.

A narrative synthesis of the extracted data was first performed to summarize the general characteristics of the included studies and their key qualitative findings. For the quantitative synthesis, a meta-analysis of prevalence was conducted for the three predefined ocular morbidities. The pooled prevalence for each outcome was calculated as a weighted average of the prevalence reported in the individual studies. Given the anticipated high degree of clinical and methodological diversity across studies, a random-effects model using the DerSimonian and Laird method was chosen a priori for all analyses. The extent of statistical heterogeneity was quantified using the I^2 statistic. I^2 values of <25%, 25%-75%, and >75% were interpreted as representing low, moderate, and high heterogeneity, respectively. The statistical significance of the heterogeneity was assessed using the Chi-squared test, with a p-value <0.10 indicating significant heterogeneity. To investigate the sources of high heterogeneity ($I^2 > 75\%$), a pre-specified sensitivity analysis was planned, wherein the meta-analysis would be re-run after systematically removing one study at a time to assess its influence on the overall pooled estimate and the I^2 value. Assessment of publication bias was performed by generating and visually inspecting a funnel plot for each outcome with sufficient studies. Asymmetry in the funnel plot can suggest the presence of publication bias. All statistical analyses were performed using Review Manager (RevMan) software, Version 5.4.

3. Results

Figure 1 showed the detailed and systematic process of study selection for the meta-analysis, presented as a PRISMA 2020 flow diagram. The process began with the Identification stage, where an initial database search yielded a substantial pool of 714 records. Before any manual screening could occur, a significant number of records were removed. The largest portion of this reduction was due to the elimination of 628 duplicate records, highlighting the overlap in database indexing. An additional 51 records were excluded by automated tools, leading to a total of 679 records being removed at this preliminary stage. This initial automated culling left 35 unique records to advance to the manual screening phase. In the subsequent Screening stage, the titles and abstracts of these 35 records were carefully reviewed against the study's core objectives. This crucial step led to the exclusion of 21 records that were deemed irrelevant to the research question. Consequently, 14 reports were considered potentially eligible and were sought for a more detailed full-text assessment. The Eligibility stage involved a rigorous multi-step assessment of these 14 reports. The first barrier was retrieval, where 2 reports could not be obtained, leaving 12 reports for full-text eligibility review. Upon detailed examination of these 12 articles, a further 6 reports were excluded for failing to meet the predefined inclusion criteria. The reasons for exclusion were clearly delineated: two were removed due to language discrepancies, two for focusing on an irrelevant type of exposure, one for being the wrong study type (a review), and one for being a case report, which is unsuitable for a meta-analysis of prevalence. Finally, the Included stage represents the culmination of this meticulous filtration process. After all exclusions, 6 studies satisfied all eligibility requirements and were carried forward for inclusion in the final quantitative synthesis, or meta-analysis. This systematic flow ensures the robustness and validity of the review's findings by clearly documenting the disposition of every record identified.

PRISMA Flow Diagram



Figure 1. PRISMA flow diagram.

Figure 2 showed a detailed summary of the key characteristics and methodological quality of the six observational studies that formed the evidence base for the meta-analysis. The table provides a transparent overview of the foundational data, allowing for a critical appraisal of the type and quality of evidence synthesized in the review. The included

studies were varied in their design, consisting entirely of non-experimental, observational research. Specifically, the evidence base comprised one descriptive cross-sectional study (Study 1), one descriptive case-series (Study 2), one retrospective case-series (Study 3), and three prospective case-series (Studies 4, 5, and 6). This highlights that the

meta-analysis draws its conclusions from real-world clinical observations rather than controlled trials. The sample sizes of the individual studies ranged considerably, from a small series of 13 patients in Study 5 to a larger cohort of 57 patients in Study 3. This variation underscores the fragmented nature of the existing literature, which consists of multiple, relatively small-scale investigations. A crucial component of the figure is the "Risk of Bias (JBI)" assessment, which evaluates the methodological rigor

of each study. The findings from this quality appraisal are particularly informative. Notably, none of the six studies were judged to have a low risk of bias, indicating that the entire evidence base possesses certain inherent methodological limitations. The assessment revealed that four of the studies (Studies 1, 3, 5, and 6) were categorized as having a moderate risk of bias. Furthermore, two studies (Studies 2 and 4) were deemed to have a high risk of bias.

Characteristics and Quality of Included Studies

STUDY ID	STUDY DESIGN	TOTAL SAMPLE (N)	RISK OF BIAS (JBI)
Study 1	Descriptive Cross-Sectional	29	Moderate
Study 2	Descriptive Case-Series	40	High
Study 3	Retrospective Case-Series	57	Moderate
Study 4	Prospective Case-Series	19	High
Study 5	Prospective Case-Series	13	Moderate
Study 6	Prospective Case-Series	21	Moderate

Figure 2. Characteristics and quality of included studies.

Figure 3 showed a comprehensive, study-by-study breakdown of the detailed patient demographics and clinical presentation findings from the six studies included in the meta-analysis. The individuals predominantly affected by these ocular injuries were young adults, with mean ages clustering in the early twenties. This was further supported by a strong and unambiguous male predominance reported across all studies that provided gender data, with male patients constituting between 69% and 87.5% of the cohorts.

This demographic signature points towards a specific at-risk population, likely reflecting the most active participants in festival celebrations. Regarding the clinical characteristics of the injuries, the data on laterality revealed that a majority of patients sustained bilateral involvement, with reported rates of 55.0%, 56.1%, and 61.9% in the studies that documented this finding. This pattern is clinically significant as it reflects the mechanism of exposure, where thrown or smeared powders inevitably contact both eyes. The

common presenting symptoms formed a classic toxidrome of acute ocular surface irritation, including burning sensations, redness, pain, watering, photophobia, and foreign body sensation. Perhaps most critically, the figure detailed the spectrum of visual acuity (VA) at presentation. While a reassuring majority of patients (between 70% and 74%) presented with Good VA (>20/40), the data also consistently

highlighted a non-trivial risk of more severe outcomes. Across three studies, a significant minority of patients suffered from Moderate VA (19-21%) or Poor VA (5-10%). This finding is crucial as it demonstrates that while most injuries may be superficial, a persistent and predictable risk of significant, vision-threatening impairment exists at the time of initial medical contact.

Detailed Patient Demographics and Clinical Presentation by Study




Figure 3. Detailed patient demographics and clinical presentation by study.

Figure 4 showed a series of three forest plots, providing a detailed quantitative synthesis of the meta-analysis for the most common ocular morbidities associated with colored powder exposure. Each plot visually and statistically summarizes the prevalence of a specific condition across the included studies. Plot A, detailing chemical conjunctivitis, revealed this to be a near-universal finding, with a pooled prevalence of 95% (95% CI: 87%-100%). The individual study estimates were consistently high, with the exception of Study 5, underscoring that an acute inflammatory response of the conjunctiva is the most immediate and frequent consequence of

exposure. However, the analysis also reported substantial heterogeneity ($I^2 = 82\%$), which is a clinically significant finding. This high value suggests that while the diagnosis of "conjunctivitis" was common, the underlying severity likely varied widely across the studies, from mild hyperemia to more severe inflammation. Plot B focused on superficial punctate keratopathy (SPK), a condition characterized by fine, diffuse erosions on the corneal surface. The pooled prevalence was a substantial 75% (95% CI: 57%-90%). The most striking feature of this analysis was the complete lack of heterogeneity ($I^2 = 0\%$). This remarkable consistency across different studies

establishes SPK as a core, pathognomonic feature of this type of ocular injury, reflecting the combined effect of chemical cytotoxicity and mechanical abrasion from the powder particles. Plot C assessed the prevalence of corneal epithelial defects, representing a more severe injury where the corneal barrier is breached. The meta-analysis found a pooled prevalence of 37% (95% CI: 23%-53%). This indicates

that while not as universal as conjunctivitis or SPK, a substantial minority of patients sustain an injury severe enough to cause a complete loss of the epithelial layer. This finding is critical as it highlights the significant risk of secondary complications, such as microbial keratitis, in over a third of affected individuals.




Figure 4. Forest plots of pooled prevalence for ocular morbidities.

Figure 5 showed the results of the meta-analysis's methodological robustness checks, presented in two distinct panels that assessed the stability of the

findings and the potential for publication bias for the chemical conjunctivitis outcome. Panel A detailed the Sensitivity Analysis, a critical procedure used to

determine if the overall results were disproportionately influenced by any single study. The table demonstrates the effect of systematically removing one study at a time and recalculating the pooled prevalence and statistical heterogeneity. The key finding was that despite the removal of any individual study, the new pooled prevalence remained high (ranging from 93% to 99%), and more importantly, the heterogeneity remained substantial, with the I^2 value consistently staying well above 75% (ranging from 79% to 88%). This powerful result indicates that the high degree of variability is not attributable to one outlier; rather, it is an inherent and consistent feature of the evidence base, likely reflecting genuine clinical diversity among the patient populations and study designs. This confirms the robustness of the primary finding that chemical conjunctivitis is a highly

prevalent issue. Panel B presented a Funnel Plot to visually assess the potential for publication bias. The plot, which maps study precision against effect size, should ideally show a symmetrical distribution of studies within the funnel. The visual inspection of this plot, however, reveals some asymmetry, with a relative lack of smaller, less precise studies showing a lower prevalence of conjunctivitis. This pattern suggests a potential publication bias, wherein studies reporting more dramatic or statistically significant findings are more likely to be published than smaller studies with null or less striking results. However, the interpretation is rightly tempered with a crucial caveat: with only four studies included in the analysis, the power to definitively detect bias is limited, and the observed asymmetry should be interpreted with caution.

Sensitivity and Publication Bias Assessment




Figure 5. Sensitivity and publication bias assessment.

4. Discussion

This systematic review and meta-analysis provide the first quantitative synthesis of ocular morbidity associated with colored powder festivals, moving the

discourse from descriptive case reports to evidence-based risk assessment.¹¹ Our analysis of 189 patients across 6 studies confirms that participation in these events carries a significant and measurable risk of

acute ocular injury. The principal finding is the remarkably high pooled prevalence of ocular surface disease: chemical conjunctivitis was present in 95% of patients, superficial punctate keratopathy in 75%, and a frank corneal epithelial defect in 37%. These findings lend robust statistical weight to the previously descriptive accounts of "Holi eye." Furthermore, our review corroborates the consistent demographic profile of affected individuals as predominantly young males between 20 and 30 years old, a finding likely reflective of the primary participants in the more boisterous aspects of these

celebrations. The high rate of bilateral involvement underscores the indiscriminate nature of the exposure mechanism. The quantitative results of this meta-analysis align with and are explained by the known pathophysiology of injury from modern synthetic powders.¹² The ocular surface, a delicate and precisely maintained biological interface, is assaulted by a combination of chemical, mechanical, and biological insults, and the clinical findings are a direct manifestation of the tissue's response to this multi-pronged attack.¹³




Figure 6. Conceptual illustration of the multi-modal ocular surface injury cascade from colored powder exposure. (A) The healthy ocular surface. (B) The initial insult combining alkaline chemical burn (liquefactive necrosis) and mechanical abrasion from particles. (C) The subsequent inflammatory cascade involves PMNs and cytokines. (D) The potential for long-term sequelae, including goblet cell loss and limbal stem cell damage.

Figure 6 showed a clear, four-part schematic that provides a scientific and narrative illustration of the multi-modal ocular surface injury cascade following exposure to colored powder. The figure effectively visualizes the progression from a healthy state to acute injury, the subsequent inflammatory response, and the potential for severe, long-term complications.

Panel A, "Healthy Ocular Surface," establishes the baseline by depicting a pristine, multi-layered epithelial barrier. This smooth and intact surface, lubricated by a healthy tear film and populated by essential goblet cells, represents the eye's normal, well-defended state. This panel underscores the biological structures that are placed at risk during

festival celebrations. Panel B, "Initial Insult," illustrates the critical moment of injury. The graphic shows colored powder particles bombarding the eye, leading to a visible epithelial defect. The accompanying text explains that this damage is a result of a "two-pronged attack": the mechanical abrasion from sharp particles physically scrapes away cells, while the alkaline chemical burn causes liquefactive necrosis, dissolving cell membranes and compromising the epithelial barrier. This stage represents the direct, acute trauma that initiates the entire pathological process. Panel C, "Inflammatory Cascade," visualizes the body's immediate biological response to this trauma. The tissue injury triggers an intense inflammatory reaction, characterized by the infiltration of immune cells, specifically polymorphonuclear leukocytes (PMNs). As depicted, these cells release a host of pro-inflammatory cytokines, such as TNF- α and IL-1 β , which amplify tissue damage and lead to the clinical signs of vasodilation (redness) and swelling. Panel D, "Long-Term Sequelae," portrays the potential chronic and devastating outcomes if the initial injury is severe and does not heal properly. The graphic shows an irregular, damaged surface with stromal scarring. The panel explains that this can include the destruction of goblet cells, leading to chronic dry eye, and damage to the vital limbal stem cells (LSCD), which can cause permanent vision impairment and chronic pain. This final panel serves as a stark warning of the permanent "hidden hazards" of these exposures.

The near-universal prevalence of chemical conjunctivitis (95%) is the most immediate and predictable outcome. It is critical to understand that this is not a simple irritation but a true chemical burn. The high alkalinity of the powders causes liquefactive necrosis of the conjunctival epithelium, triggering a massive inflammatory cascade mediated by histamine, IL-8, and TNF- α , resulting in the clinically observed redness and swelling.¹⁴ The high statistical heterogeneity ($I^2=82\%$) in this finding is not surprising and is clinically informative. It likely reflects the broad spectrum of conjunctival injury, from mild, transient

hyperemia to severe pseudomembrane formation, all of which would be classified under the umbrella of "conjunctivitis" in the primary studies. Simultaneously, the high prevalence of superficial punctate keratopathy (75%) reflects a more specific injury to the cornea. This pattern of fine, scattered epithelial erosions arises from the synergistic mechanisms of direct cytotoxicity from the chemical dyes, which induce apoptosis via oxidative stress, and the mechanical abrasive effect of silica and mica particles. The remarkable consistency between studies for this finding ($I^2=0\%$) suggests that SPK is a core, pathognomonic feature of this type of injury.¹⁵

Our finding that a substantial proportion develop a frank corneal epithelial defect (37%) represents a clinically significant escalation of injury. This breach of the corneal barrier is the critical gateway to more severe complications. The clinical importance of an epithelial defect cannot be overstated, as it creates a high-risk scenario for secondary infectious keratitis. However, the "hidden hazard" of these celebrations extends far beyond the acute phase. Even after the initial injury appears to heal, the chemical insult can initiate a lifetime of ocular surface disease. This is a crucial aspect often overlooked in acute care settings. The alkaline powders and cytotoxic dyes can destroy the mucin-producing goblet cells in the conjunctiva. The loss of this crucial tear film layer leads to evaporative dry eye, chronic foreign body sensation, photophobia, and a lifelong dependency on artificial tears and other therapies. The limbus, at the periphery of the cornea, houses the stem cells responsible for replenishing the corneal epithelium. A severe chemical burn can destroy these stem cells. The consequence is limbal stem cell deficiency (LSCD), a devastating condition where the cornea becomes covered by opaque, vascularized conjunctival tissue, leading to chronic pain, recurrent erosions, and severe, often irreversible, vision loss.¹⁵ Damage to the epithelial basement membrane and its anchoring fibrils (Type VII collagen) can lead to recurrent corneal erosion syndrome (RCES), a condition where the corneal epithelium fails to adhere properly and

spontaneously sloughs off, causing episodes of excruciating pain, typically upon waking. Even in the absence of a dense central scar, the ocular surface damage can permanently degrade the quality of vision. Stromal haze, basement membrane irregularities, and chronic inflammation can induce significantly higher-order aberrations, leading to debilitating symptoms of glare, halos, starbursts, and reduced contrast sensitivity.¹⁶ A patient may read the 20/20 line on a chart but be functionally blind when driving at night.

The findings of this study provide a clear mandate for public health action. While the data is derived from the Indian Holi festival, the pathophysiological principles are universal, serving as a critical warning for any festival involving unregulated powders. The recommendations must be pragmatic and multifaceted. The cornerstone of clinical advice is often the pursuit of an ideal standard of care. In the context of preventing ocular trauma from colored powder, the unequivocal ideal is the use of sealed, wrap-around safety goggles.¹⁷ This Gold Standard represents the ophthalmologist's choice, providing a near-impermeable barrier that protects the ocular surface not only from the direct, forceful impact of thrown powder clumps but also from the fine, aerosolized dust that permeates the celebratory atmosphere. This complete seal is critical because it mitigates both the mechanical and chemical insults simultaneously. The physical barrier prevents the sharp, abrasive particles of silica or mica from causing corneal erosions, while the enclosed space protects the delicate tear film from being saturated with alkaline chemicals that can rapidly destabilize its pH and initiate a cascade of tissue damage. In essence, sealed goggles offer a personal "clean room" environment for the eyes in the midst of a hazardous particulate storm. However, we must rigorously acknowledge the significant behavioral and social barriers to the adoption of this Gold Standard. For many participants, particularly young travelers, the primary motivation for attending such festivals is the pursuit of an immersive, authentic, and sensory experience. The act of wearing bulky safety goggles can feel antithetical to this goal.

They can be perceived as hot, uncomfortable, and visually limiting, but most importantly, they can create a sense of detachment, transforming an active participant into a mere observer. This social-psychological barrier is immensely powerful and often leads to low compliance, rendering the perfect-in-theory advice useless in practice.

Therefore, a pragmatic public health message must incorporate a Good Alternative based on the principles of harm reduction. This alternative is the use of well-fitting, high-coverage sunglasses. While not sealed, a wrap-around style of sunglass can deflect the vast majority of direct projectile impacts. The physics of this intervention is simple but effective: by presenting a physical barrier, sunglasses prevent the largest and most dangerous clumps of powder from making direct contact with the ocular surface. This intervention is designed to significantly reduce the total load of the hazardous material reaching the eye. The difference in the total volume of powder exposure can be the critical factor that distinguishes a mild, transient irritation from a severe, vision-threatening chemical burn requiring prolonged medical care.¹⁸ The key attributes are "well-fitting" and "high-coverage," ensuring that the gap between the frame and the face is minimized, thereby offering a degree of protection from particles entering from the sides, top, or bottom. This approach represents a realistic compromise, offering substantial protection while being far more socially acceptable and comfortable for the user.

Finally, for individuals who may reject even sunglasses, a Minimum Standard of protection must be communicated. This final safety net emphasizes that any physical barrier is superior to none. Even standard prescription eyeglasses can block a direct impact that might otherwise have devastating consequences. This equipment-based advice is then coupled with a crucial emphasis on behavioral modification. This is not a passive approach but an active form of self-protection that requires situational awareness. Participants should be counseled to be mindful of their surroundings, to anticipate when powder is about to be thrown, and to instinctively turn

their head or use their hands to shield their face. The simple, reflexive act of closing the eyes tightly just before an anticipated exposure can prevent the vast majority of powder from reaching the conjunctiva and cornea. This tiered strategy—from the ideal to the pragmatic to the minimal—allows a health advisor to provide tailored, realistic advice. It moves away from a single, often-rejected directive and towards a collaborative approach that respects individual choice while maximizing the chances that some form of protective measure will be adopted, ultimately reducing the overall incidence and severity of injury.

While empowering individuals with harm reduction strategies is essential, placing the entire burden of safety on the participant is an incomplete and fundamentally flawed public health model. A truly effective prevention strategy must adopt a "systems thinking" approach, modifying the environment in which the event takes place and addressing the root cause of the hazard—the composition of the powders themselves. This requires a shift in focus from solely managing individual behavior to engineering a safer ecosystem for the celebration. The most immediate and high-impact environmental modification is the implementation of event-level safety measures, specifically the establishment of dedicated irrigation stations. This intervention acts as a form of immediate post-exposure prophylaxis. These stations should be clearly marked with universally recognizable symbols, strategically placed throughout the event venue, and staffed with personnel trained in basic ocular first aid. They should be equipped with large, accessible sources of sterile, buffered saline solution or, at a minimum, an abundant supply of clean, bottled water. The scientific rationale for this is clear and compelling. The primary mechanism of injury from these powders is an alkali chemical burn. Immediate, copious, and prolonged irrigation is the single most effective intervention to mitigate this damage.¹⁹ Irrigation works by physically washing away foreign particles, diluting the concentration of harmful chemicals, and, most importantly, neutralizing the pH of the ocular surface, halting the process of liquefactive necrosis

before it can cause deeper and more permanent tissue damage. The presence of such stations transforms the safety paradigm from one of solely pre-event prevention to one that includes a robust, on-site emergency response, dramatically improving outcomes for those who do sustain an exposure despite their best efforts. Moving further upstream to address the root of the problem requires interventions at the supply chain and vendor regulation level. The fundamental issue is not the cultural practice of celebrating with color, but the hazardous nature of the modern, synthetic materials used. A powerful public health strategy would therefore focus on shifting the market back towards safe, traditional, and non-toxic powders. This can be achieved through a multi-pronged campaign in endemic regions. First, public awareness campaigns are needed to educate local communities and vendors about the documented health dangers of industrial dyes, linking them to specific outcomes like eye injuries and skin allergies. This messaging should be paired with a positive campaign that celebrates and promotes the use of authentic, natural *gulal* derived from traditional sources like turmeric, beetroot, and flower extracts, framing the choice as one that is not only safer but also more culturally authentic.

These campaigns work to create consumer demand for safer products. When participants begin to preferentially purchase powders that are certified as non-toxic or are visibly organic, vendors are economically incentivized to alter their supply chains. This market-based approach can be powerfully supplemented by supporting local producers and artisans who create safe, traditional powders. By creating a positive economic feedback loop, public health initiatives can make the safe choice the easy and economically beneficial choice for the entire community. This upstream intervention is the most sustainable solution, as it removes the hazard at its source rather than simply attempting to manage its consequences. Finally, any discussion of public health interventions must maintain a Balanced Focus, acknowledging the crucial importance of equity. While

the implications for travel medicine are a novel and important focus, it is essential to recognize that the overwhelming burden of injury is borne by the local residents who participate in these festivals annually, often over a lifetime. A public health strategy that focuses exclusively on making an event safe for tourists while ignoring the endemic risks to the host community is ethically untenable.²⁰ Therefore, the most effective and equitable interventions are those designed with and for the local community. System-level changes like vendor regulation and the promotion of safe powders primarily benefit local residents, with the wonderful secondary effect of creating a safer environment for visitors. Travel medicine advice then becomes a complementary layer of personal protection within a system that is fundamentally safer for everyone. The ultimate goal should be to foster a celebratory environment where the well-being of the local population and international travelers are seen not as separate issues, but as inextricably linked components of a single, healthy community.

5. Conclusion

This meta-analysis demonstrates that participation in colored powder festivals, based on extensive evidence from Holi in India, carries a substantial and quantifiable risk of acute ocular morbidity, including exceptionally high rates of chemical conjunctivitis and corneal epithelial damage. The findings transition the understanding of these injuries from a series of isolated incidents to a predictable and preventable public health problem. The "hidden hazard" of celebration extends beyond the acute injury to a lifetime of potential ocular surface disease. It is therefore imperative that national public health bodies and international travel medicine organizations incorporate these findings into their guidelines, advocating for pragmatic, tiered preventive strategies and educating both local participants and international tourists to ensure that these vibrant celebrations of life and color do not result in avoidable, and potentially irreversible, harm to sight.

6. References

1. Wagh V, Tidake P. Clinical study and profile of ocular trauma: Findings from a rural hospital in central India. *Cureus*. 2022; 14(7): e26915.
2. Sanjay S, Gupta A, Uday S. Scare air gun-induced ocular trauma in an agriculture farm in India. *Pan-Am J Ophthalmol*. 2022; 4(1): 56.
3. Chauhan R, Singh A, Mishra S, Baranwal VK. Epidemiological profile of ocular trauma in A mid-zonal hospital of armed forces in North-East region of India. *Delhi J Ophthalmol*. 2021; 31(4): 63–6.
4. Wadwekar B, Mishra AK, Sahu P, Sanjana E, Manikandan M. Burden of ocular trauma and its effect on vision. *J Curr Res Sci Med*. 2021; 7(2): 97–101.
5. Bhat K, Gupta K, Khandelwal R. Ocular trauma study in rural Indian population – clinical features and visual outcomes based on ocular trauma score. *J Clin Ophthalmol Res*. 2024; 12(2): 154–8.
6. Rana A, Kumar J, Goynka Y. Clinical study and profile of ocular trauma: Findings from a tertiary care centre from northern India. *Int J Adv Res (Indore)*. 2025; 13(06): 1211–5.
7. Sharma B, Singh S, Kumar K, Dubey A, Ahirwar K. Epidemiology, clinical profile and factors, predicting final visual outcome of pediatric ocular trauma in a tertiary eye care center of Central India. *Indian J Ophthalmol*. 2017; 65(11): 1192.
8. Qayum S, Anjum R, Rather S. Epidemiological profile of pediatric ocular trauma in a tertiary hospital of northern India. *Chin J Traumatol*. 2018; 21(2): 100–3.
9. Madanagopalan VG, Nagesha CK, Velis G, Devulapally S, Balamurugan S. Frosted branch angiitis with penetrating ocular trauma and retained intraocular foreign body. *Indian J Ophthalmol*. 2018; 66(7): 1031–3.

10. Mishra S, Goel S, Nangia P, Senger D, Shah AV, Saurabh K, et al. Acute macular neuroretinopathy after blunt ocular trauma: a rare association. *Indian J Ophthalmol.* 2019; 67(4): 566–8.
11. Bhole PK, Mansuri F, Parmar D, Gandhi K. Impact of COVID-19 pandemic on the spectrum of ocular trauma during Diwali at a tertiary eye care center of Western India. *Indian J Ophthalmol.* 2021; 69(12): 3789–90.
12. Bhatti A, Kakkar A, Singh S. Profile of ocular injuries amongst various trauma patients presenting to tertiary care centre of Punjab. *Indian J Clin Exp Ophthalmol.* 2022; 7(4): 642–7.
13. Singh A, Maurya RP, Singh VP, Gautam S, Kumar A, Mishra CP, et al. Study of epidemiology, clinical profile, visual outcome and prognostic factors of blunt ocular trauma in a teaching hospital. *Indian J Clin Exp Ophthalmol.* 2022; 8(4): 458–67.
14. Vaghmare S, Rahul R, Prajna NV, Radhakrishnan N. Analysis on the completeness of case records of patients with penetrating ocular trauma following a multimodal intervention. *Indian J Ophthalmol.* 2022; 70(8): 2962–5.
15. Wadwekar B, Hansdak A, Sahu P, Karnam AHF, Sanjana EF. Clinical profile of ocular trauma at a tertiary care hospital in South India: a retrospective study. *Trauma.* 2023; 25(1): 48–52.
16. Goswami K, Mazumdar I. A study of the levels of some toxic substances present in dry Holi colours in Kolkata, India. *Indian J Clin Biochem.* 2022; 37(2): 206–11.
17. Kukfisz B, Piec R. The fire and explosion hazard of coloured powders used during the Holi festival. *Int J Environ Res Public Health.* 2021; 18(21): 11090.
18. Gupta S, Selvan H, Markan A, Gupta V. Holi colors and chemical contact keratitis. *EYE.* 2018; 32(1): 1–3.
19. Pujari A, Behera A, Mukhija R, Chawla R, Yadav S, Sharma N. Ocular toxicity due to colours used during Holi celebration in India: correlation of clinical findings with the anterior segment OCT. *Cutan Ocul Toxicol.* 2019; 38(1): 1–4.
20. Garg N, Gupta SK, Bhatnagar A, Hanumanthappa V. A study on after effects of Holi: Color injury, ocular trauma, and assault at a tertiary care ophthalmic center in north India. *Int J Res Found Hosp Healthc Adm.* 2018; 6(1): 31–6.